arXiv:1508.05364v1 [math.CO] 21 Aug 2015

K-CIRCULAR MATROIDS OF GRAPHS

José F. De Jesús

University of Puerto Rico, San Juan, Puerto Rico, United States

Alexander Kelmans

University of Puerto Rico, San Juan, Puerto Rico, United States
Rutgers University, New Brunswick, New Jersey, United States

Abstract

In 30's Hassler Whitney considered and completely solved the problem (WP) of describing the classes of graphs G having the same cycle matroid $M(G)$ [15, 16]. A natural analog $(WP)'$ of Whitney's problem (WP) is to describe the classes of graphs G having the same matroid $M'(G)$, where $M'(G)$ is a matroid (on the edge set of G) distinct from $M(G)$. For example, the corresponding problem $(WP)' = (WP)_\theta$ for the so-called bicircular matroid $M_\theta(G)$ of graph G was solved in [2, 11]. We define the so-called *k-circular matroid* $M_k(G)$ on the edge set of graph G for any non-negative integer k so that $M(G) = M_0(G)$ and $M_\theta(G) = M_1(G)$. It is natural to consider the corresponding analog $(WP)_k$ of Whitney's problem (WP) not only for $k = 0$ and $k = 1$ but also for any integer $k \geq 2$. In this paper we give a characterization of the k-circular matroid $M_k(G)$ by describing the main constituents (circuits, bases, and cocircuits) in terms of graph G and establish some important properties of the k-circular matroid. The results of this paper will be used in our further research on the problem $(WP)_k$. In our next paper we use these results to study a particular problem of $(WP)_k$ on graphs uniquely defined by their k-circular matroids.

Key words: graph, bicycle, cacti-graph, splitter theorems, matroid, k-circular matroid.

MSC Subject Classification: 05B35, 05C99

1 Introduction

In 30's Hassler Whitney developed a remarkable theory on the matroid isomorphism and the matroid duality of graphs [14–17]. He considered a graph G and the so called cycle matroid $M(G)$ of G (whose circuits are the edge subsets of the cycles in G) and stated the following natural problems on pairs $\langle G, M(G)\rangle$:

(WP) describe the classes of graphs having the same cycle matroid and, in particular, graphs that can be reconstructed from cycle matroid (up to the names of vertices) and

(WP^*) describe the pairs of graphs whose cycle matroids are dual, i.e. describe the class of graphs closed under their cycle matroids duality.

Classical Whitney's graph matroid-isomorphism theorem and Whitney's planarity criterion provide the answers to the above questions [15–17] (see also [10]).

Naturally, Whitney's problems and interesting results along this line prompted further questions and research on possible strengthenings as well as various extensions or analogs of some Whitney's results (see, for example, [2, 5–7, 9, 11]).

The goal of this research is twofold:

($\Gamma 1$) to introduce and consider some new matroids related with a graph G that are distinct from the cycle matroid $M(G)$ and to establish the structural properties of those matroids in terms of G and

($\Gamma 2$) to extend some of Whitney's results on problem (WP) to this variety of graph matroids using their structural properties established in ($\Gamma 1$)

This paper is the first one in the series of our papers along the ($\Gamma 1$) - ($\Gamma 2$) line on graphs and their matroids.

Section 2 provides some basic notions, notation, and some known facts on matroids and graphs.

In Section 3 some additional notions and some auxiliary and preliminary facts on graphs are presented that will be used later. In particular, we will describe some useful properties of the function $\Delta(G) = |E(G)| - |V(G)|$ of graph G, establish the splitter theorems for graphs with no leaves and no cycle components. We also define and characterize two special subgraphs of a graph that are called the *core* and the *kernel* of a graph. These notions and results will play an essential role in the study of the so-called k-circular matroids.

In Section 4 we introduce the notion of the k-circular matroid $M_k(G)$ of a graph G, where k is a non-negative integer, and establish some properties of this matroid. In particular, we describe the main constituents of this matroid (bases, circuits, cocircuits, etc.) in graph terms.

The results of this paper provide, in particular, a proper basis for our study of the problem $(WP)_k$ on describing the classes of graphs with the same k-circular matroids.

2 Basic notions and facts on matroids and graphs

2.1 On clutter and hereditary families

Given a partial order set $(P, \preceq)$, a maximal element of $(P, \preceq)$ is called $\preceq$-*maximal* and a minimal element of $(P, \preceq)$ is called $\preceq$-*minimal*.

The notions and facts described in this Section can be found in [8]. Given a finite set E, consider the poset $\mathcal{P} = (2^E, \subseteq)$ and $\mathcal{X} \subseteq 2^E$.

Let
$\mathcal{M}ax(\mathcal{X})$ denote the set of $\subseteq$-maximal elements of poset $(\mathcal{X}, \subseteq)$ and
$\mathcal{M}in(\mathcal{X})$ denote the set of $\subseteq$-minimal elements of poset $(\mathcal{X}, \subseteq)$.

Let
$\mathcal{M}ax^{-1}(\mathcal{X}) = \{X \subseteq E : X \subseteq Y \ for\ some\ Y \in \ \mathcal{X}\}$ and
$\mathcal{M}in^{-1}(\mathcal{X}) = \{X \subseteq E : X \supseteq Y \ for\ some\ Y \in \ \mathcal{X}\}$.

A family $\mathcal{X}$ is called a *clutter* if $\mathcal{M}ax(\mathcal{X}) = \mathcal{M}in(\mathcal{X})$, i.e. if $X \not\subseteq Y$ for every $X, Y \in \mathcal{X}$ and $X \neq Y$. Obviously, $\mathcal{M}ax(\mathcal{X})$ and $\mathcal{M}in(\mathcal{X})$ are clutters.

A family $\mathcal{X}$ is called *hereditary* if $X \subseteq Y \in \mathcal{X} \Rightarrow X \in \mathcal{X}$.

A family $\mathcal{X}$ is called *anti–hereditary* if $X \supseteq Y \in \mathcal{X} \Rightarrow X \in \mathcal{X}$.

Obviously, the following is true.

Claim 2.1.1 *Let* $\mathcal{X} \subseteq 2^E$. *Then*
$(c1)$ $\mathcal{M}ax^{-1}(\mathcal{X})$ *is a hereditary family,*
$(c2)$ $\mathcal{M}in^{-1}(\mathcal{X})$ *is an anti-hereditary family,*
$(c3)$ $\mathcal{X}$ *is a hereditary family if and only if* $\mathcal{M}in^{-1}(\mathcal{M}ax(\mathcal{X})) = \mathcal{X}$, *and*
$(c4)$ $\mathcal{X}$ *is an anti-hereditary family if and only if* $\mathcal{M}ax(\mathcal{M}in^{-1}(\mathcal{X})) = \mathcal{X}$.

Let $\mathcal{I} \subseteq 2^E$ and $\mathcal{I}$ a hereditary family and put $H = (E, \mathcal{I})$ and $\mathcal{I} = \mathcal{I}(H)$. An element X of $\mathcal{I}$ is called an *independent set of* H and $\mathcal{I}$ is the *independence family of* H.

Let $\mathcal{B} = \mathcal{B}(H) = \mathcal{M}ax(\mathcal{I})$. An element B of $\mathcal{B}$ is called a *base of* H and $\mathcal{B} = \mathcal{B}(H)$ is the *family of bases of* H.

Let $\mathcal{D} = 2^E \backslash \mathcal{I}$. Obviously, $D \in \mathcal{D}$ if and only if X is not independent set of H. Therefore an element D of $\mathcal{D}$ is called a *dependent set of* H and $\mathcal{D} = \mathcal{D}(H)$ is the *dependence family of* H.

Let $\mathcal{C} = \mathcal{M}in(\mathcal{D})$. An element C of $\mathcal{C}$ is called a *circuit of* H and $\mathcal{C} = \mathcal{C}(H)$ is the *family of circuits of* H. A circuit C of H consisting of one element is called a *loop of* H.

Given $\mathcal{B} = \mathcal{B}(H)$, let $\mathcal{B}^* = \mathcal{B}^*(H) = \{E \setminus B : B \in \mathcal{B}\}$, $\mathcal{I}^* = \mathcal{M}ax^{-1}(\mathcal{B}^*)$, $\mathcal{D}^* = 2^E \setminus \mathcal{I}^*$, $\mathcal{C}^* = \mathcal{M}in(\mathcal{D}^*)$, and $H^* = (E, \mathcal{I}^*)$. We call $H = (E, \mathcal{I})$ and $H^* = (E, \mathcal{I}^*)$ *dual hereditary families on* E and put $\mathcal{B}(H^*) = \mathcal{B}^*(H)$, $\mathcal{D}(H^*) = \mathcal{D}^*(H)$, and $\mathcal{C}(H^*) = \mathcal{C}^*(H)$.

Accordingly, we call
an independent set of H^* a *coindependet set of* $H = (E, \mathcal{I})$,
a dependent set H^* a *codependent set of* $H = (E, \mathcal{I})$,
a base of H^* a *cobase of* $H = (E, \mathcal{I})$,
a circuit of H^* a *cocircuit of* $H = (E, \mathcal{I})$, and
a loop of H^* a *coloop of* $H = (E, \mathcal{I})$.

It is easy to see that the following is true.

Claim 2.1.2 *Let* $\mathcal{I} \subseteq 2^E$, *and* $\mathcal{I}$ *is a hereditary family. Let* $H = (E, \mathcal{I})$ *and* $\mathcal{I} = \mathcal{I}(H)$. *Then*
$(c0)$ $\mathcal{I}^*(H)$ *is a hereditary family,*
$(c1)$ $\mathcal{B}(H)$ *and* $\mathcal{B}^*(H)$ *are clutters,*
$(c2)$ $\mathcal{D}(H)$ *and* $\mathcal{D}^*(H)$ *are anti-hereditary families,*

($c3$) $\mathcal{C}(H)$ and $\mathcal{C}^*(H)$ are clutters, and

($c4$) every family in $\{\mathcal{I}(H),\ \mathcal{B}(H),\ \mathcal{D}(H),\ \mathcal{C}(H),\ \mathcal{I}^*(H),\ \mathcal{B}^*(H),\ \mathcal{D}^*(H),\ \mathcal{C}^*(H)\}$ is uniquely defined by any other family in the above list.

It is also easy to prove the following.

Claim 2.1.3 *Let $H = (E, \mathcal{I})$ be a non-empty hereditary family of subsets of E. Then*

($c1$) $B \in \mathcal{B}(H)$ and $C \in \mathcal{C}(H) \Rightarrow C \not\subseteq B$ or, equivalentlly,

($c2$) $B \in \mathcal{B}(H)$ and $C^* \in \mathcal{C}(H) \Rightarrow B \cap C^* \neq \emptyset$ and
$B^* \in \mathcal{B}^*(H)$ and $C \in \mathcal{C}(H) \Rightarrow B^* \cap C \neq \emptyset$.

2.2 On matroids

In this Section we describe some notions and necessary facts on matroids. Most of them can be found in [10, 12]. Given a set $X \subseteq E$ and $e \in E$, we usually write $X \cup e$ instead of $X \cup \{e\}$ and $X \setminus e$ instead of $X \setminus \{e\}$.

A *matroid* is a pair $M = (E, \mathcal{I})$, where E is a finite non-empty set and $\mathcal{I} \subseteq 2^E$ such that

($A\mathcal{I}1$) $\emptyset \in \mathcal{I}$,

($A\mathcal{I}2$) $X \subseteq Y \in \mathcal{I} \Rightarrow X \in \mathcal{I}$, and

($A\mathcal{I}3$) $X, Y \in \mathcal{I}$ *and* $|X| < |Y| \Rightarrow \exists\ y \in Y \setminus X$ *such that* $X \cup y \in \mathcal{I}$.

A set $I \in \mathcal{I}$ is called an *independent set of M*.

Notice that ($A\mathcal{I}1$) and ($A\mathcal{I}2$) imply that $\mathcal{I}$ is a hereditary family (see Section 2.1). Therefore $M = (E, \mathcal{I})$ is a matroid if and only if $\mathcal{I}$ is a non-empty hereditary family satisfying ($A\mathcal{I}3$). Thus, all notions and claims about hereditary families in Section 2.1 are valid for the matroids.

The next two claims provide criteria for a family $\mathcal{F} \subseteq 2^E$ to be the family of bases and the family of circuits of a matroid, respectively.

Claim 2.2.1 *Let $\mathcal{B} \subseteq 2^E$. Then $\mathcal{B}$ is the family of bases of a matroid if and only if*

($A\mathcal{B}1$) $\mathcal{B}$ is a non-empty clutter and

($A\mathcal{B}2$) $X, Y \in \mathcal{B}$ and $X \neq Y \Rightarrow \forall\ x \in X \setminus Y\ \exists\ y \in Y \setminus X$ such that $(X \setminus x) \cup y \in \mathcal{B}$.

From Claim 2.2.1 it follows that every two bases of matroid M are of the same size. Put $\rho(M) = |B|$ for $B \in \mathcal{B}(M)$ and $\rho^*(M) = |E| - \rho(M)$. We call $\rho(M)$ the *rank of matroid M* and $\rho^*(M)$ the *corank of matroid M*.

Claim 2.2.2 *Let $\mathcal{C} \subseteq 2^E$. Then $\mathcal{C}$ is the family of circuits of a matroid if and only if*

($A\mathcal{C}1$) $\mathcal{C}$ is a clutter such that $\emptyset \notin \mathcal{C}$ and

($A\mathcal{C}2$) $X, Y \in \mathcal{C}$ and $X \cap Y \neq \emptyset \Rightarrow \forall\ a \in X \cap Y\ \exists\ C \in \mathcal{C}$ such that $C \subseteq (X \cup Y) \setminus a$
(see Figure 1).

We will need the following simple facts.



Figure 1: Axiom $(A\mathcal{C}2)$ of a matroid

Claim 2.2.3 *Let $M = (E, \mathcal{I})$ be a matroid. Then the following are equivalent:*

$(a1)$ c is a coloop of M,

$(a2)$ $c \in B$ for every $B \in \mathcal{B}(M)$, and

$(a3)$ $c \notin C$ for every $C \in \mathcal{C}(M)$.

Claim 2.2.4 *Let $M = (E, \mathcal{I})$ be a matroid. Then*

$(c1)$ if $B \in \mathcal{B}(M)$ and $e \in E \setminus B = B^*$, then there exists a unique circuit $C = C(e, B)$ of M such that $e \in C \subseteq B \cup e$ (or, equivalently, such that $C \cap B^* = \{e\}$) and similarly, if $B \in \mathcal{B}(M)$ and $e \in B$, then there exists a unique cocircuit $C^* = C^*(e, B)$ of M such that $e \in C^* \subseteq B^* \cup e$ (or, equivalently, such that $C^* \cap B = \{e\}$), $(c2)$ $u \in C(e, B) \setminus e \Leftrightarrow (B \setminus u) \cup e \in \mathcal{B}(M)$ and similarly, $u \in C^*(e, B) \setminus e \Leftrightarrow (B \setminus e) \cup u \in \mathcal{B}(M)$,

$(c3)$ for every $C \in \mathcal{C}(M)$ $(C^* \in \mathcal{C}(M))$ there exists $B \in \mathcal{B}$ and $e \in E \setminus B = B^*$ such that $C = C(e, B)$ (respectively, $e \in B$ such that $C^* = C^*(e, B)$).

As in Section 2.1, given a matroid $M = (E, \mathcal{I})$, let $M^* = (E, \mathcal{I}^*)$ be the pair with $\mathcal{B}^*(M) = \{E \setminus B : B \in B(M)\}$. It is easy to see that M^* is a matroid and M^* is the dual of M.

Given $B \in \mathcal{B}(M)$ and $e \in E \setminus B = B^*$, we call $C(e, B)$ the *fundamental circuit of B in M rooted at e* or simply, the *(B, e)-circuit in M*.

Similarly, if $B^* \in \mathcal{B}(M^*)$ and $e \in E \setminus B^* = B$, we call $C(e, B^*)$ the *fundamental circuit of B^* in M^* rooted at e* or simply, the *(B^*, e)-circuit in M^**.

We also call $C(e, B^*)$ the *fundamental cocircuit of base B in M rooted at e*, or simply, the *(B, e)-cocircuit in M*.

Claim 2.2.5 *Let $M = (E, \mathcal{I})$ be a matroid and $a, b \in E$, $a \neq b$. Then the following are equivalent:*

$(a1)$ a and b belong to a circuit of M and

$(a2)$ a and b belong to a cocircuit of M.

Let $M = (E, \mathcal{I})$ be a matroid, L the set of loops, and L^* the set of coloops of M. Given elements a and b in $E \setminus (L \cup L^*)$ we write $a \sim b$ if a and b belong to a common circuit of M or, equivalently (by the above Claim), if a and b belong to a common cocircuit of M.

Claim 2.2.6 *$(E \setminus (L \cup L^*), \sim)$ is an equivalence relation.*

Claim 2.2.7 *Let $M = (E, \mathcal{I})$ be a matroid, $Z \subseteq E$, and $M|_Z = (Z, \{X \subseteq Z : X \in \mathcal{I}\})$. Then $M|_Z$ is a matroid.*

For $X \subseteq E$ let $M \setminus X = M|_{E \setminus X}$ and $M/X = (M^* \setminus X)^*$.

A matroid N is called a *component of matroid* $M = (E, \mathcal{I})$ if $N = M|_Z$ for some equivalence class Z of the equivalence relation $(E \setminus (L \cup L^*), \sim)$, of matroid M, and so N has at least two elements. Let $Cmp(M)$ denote the set of components of M.

A matroid $M = (E, \mathcal{I})$ is called *connected* if $|E| \geq 2$ and $a \sim b$ for every $a, b \in E$, i.e. matroid $M = (E, \mathcal{I})$ is connected if and only if M has exactly one components and M has no loops and no coloops.

Claim 2.2.8 *Matroid N is a component of matroid M if and only if N is a component of M^*, and so $Cmp(M) = Cmp(M^*)$.*

We call a circuit C of a connected matroid M a *non-separating circuit* of M if M/C is a connected matroid. Similarly, we call a cocircuit C^* of a connected matroid M a *non-separating cocircuit* of M if $M \setminus C^*$ is a connected matroid or, equivalently, if C^* is a non-separating circuit of M^*.

We call a matroid $M = (E, \mathcal{I})$ *non-trivial* if E is neither a base of M and nor a base of M^* and *trivial*, otherwise. Obviously, M is non-trivial if and only if $\mathcal{C}(M) \neq \emptyset$ and $\mathcal{C}^*(M) \neq \emptyset$.

Let $M = (E, \mathcal{I})$ and $M' = (E', \mathcal{I}')$ be matroids. An *isomorphism from M to M'* is a bijection ε from E to E' such that $A \in \mathcal{I} \Leftrightarrow \varepsilon[A] \in \mathcal{I}$, where $\varepsilon[A] = \{\varepsilon(e) : e \in A\}$. *Matroids M and M' are isomorphic* if there exists an isomorphism from M to M' (or, equivalently, an isomorphism from M' to M).

2.3 On graphs

In this Section we describe some notions and necessary facts on graphs. Most of them can be found in [1,3,13].

A *graph* G is a triple (V, E, ϕ) such that V and E are disjoint finite sets, $V \cap E = \emptyset$, $V \neq \emptyset$, and $\phi : E \to \binom{V}{2} \cup V$. The elements of $V = V(G)$ and $E = E(G)$ are called *vertices and edges of graph* G, respectively. If $\phi(e) = (v, v)$ for some $v \in V$, then e is called a *loop of* G. If $\phi(a) = \phi(b)$ for some $a, b \in E$, then a and b are called *parallel edges of* G.

Graphs $G = (V, E, \phi)$ and $G' = (V', E', \phi')$ are *equal* if $V = V'$, $E = E'$, and $\phi = \phi'$ (see Figure 2).

For graphs $G_1 = (V_1, E_1, \phi_1)$ and $G_2 = (V_2, E_2, \phi_2)$ with $E_1 \cap E_2 = \emptyset$, let G be the graph (V, E, ϕ), where $V = V_1 \cup V_2$, $E = E_1 \cup E_2$, and for $e \in E$, $\phi(e) = \phi_i(e)$ if $e \in E_i$, $i = 1, 2$. Then G is called the *union* of G_1 and G_2, written $G_1 \cup G_2$.

We called a graph $G = (V, E, \phi)$ *simple* if $\phi : E \to \binom{V}{2} \cup V$ is an injective function, and so G ha no parallel edges.



$$\boldsymbol{V(A) = V(B),\; E(A) = E(B),\; \phi_A \neq \phi_B \Rightarrow A \neq B}$$

Figure 2: Two different but isomorphic graphs

A graph (V, E, ϕ) is called *complete* if $\phi : E \to \binom{V}{2}$ is a bijection.

If $\phi(e) = (u, v)$ in a graph G, then we say that vertices u and v are *adjacent*, edge e and vertex v are *incident in* G, and u, v are the *end vertices of edge* e *in* G.

A vertex in G incident to no edge is called an *isolated vertex of* G.

A vertex v in G is called an *leaf of* G if v is incident to exactly one edge of G and this edge is not a loop.

Unless stated otherwise, we assume that a graph has no isolated vertices.

Given a graph $G = (V, E, \phi)$, an element $a \in V \cup E$, and a set $A \subseteq V \cup E$, we say that a *is incident to* A if a is incident to an element of A.

Given G and $v \in V(G)$, the v*-star* (or a *vertex star* of G) is the set $S(v, G)$ of edges in G incident to vertex v. Let $\mathcal{S}(G) = \{S(v, G) : v \in V(G)\}$ (see Figure 7).

The *degree* $d(v, G)$ *of vertex* v *in* G is the number of non-loop edges in $S(v, G)$ plus 2-times the number of loops in $S(v, G)$. The function $d : V \to \mathbb{N}_0$ such that $d(v) = d(v, G)$ for $v \in V$ is called the *degree function of* G and denoted by $d(G)$.

Let $G = (V, E, \phi)$ be a graph, $X \subset V$, and $Y \subseteq E$. Then $G \setminus X$ denotes the graph $G' = (V', E', \phi')$ such that $V' = V \setminus X$, E' is obtained from E by deleting all edges of G incident to at least one vertex in X , and $\phi' : E' \to \binom{V'}{2} \cup V'$ is the restriction of function ϕ on E'. We say that G' *is obtained from* G *by deleting vertex set* X.

Similarly, $G \setminus Y$ denotes the graph $G' = (V', E', \phi')$ such that $V' = V$, $E' = E \setminus Y$, and $\phi' : E' \to \binom{V'}{2} \cup V'$ is the restriction of ϕ on E'. We say that G' *is obtained from* G *by deleting edge set* Y.

A graph G' is a *subgraph of* G, written as $G' \subseteq^s G$, if G' can be obtained from G by deleting some edge subset Y and some vertex subset X of G. Obviously, $\subseteq^s$ is a partial order on the set of graphs.

Given a graph $G = (V, E, \phi)$ and $Y \subseteq E$, let $G\langle Y \rangle$ denote the subgraph of G obtained from G by deleting all edges in $E \setminus Y$ and all vertices of G that are incident to no edge in Y. We call $G\langle Y \rangle$ the *subgraph of G induced by Y*. We also say that Y *spans $X \subseteq V$ in G* if $X = V(G\langle Y \rangle)$. Obviously, if A is a subgraph of G with no isolated vertices, then $G\langle E(A) \rangle = A$.

Let $\mathcal{G}_2^{1,1}$ denote the set of graphs having exactly two vertices, say x and y, of degree 1 and all other vertices of degree 2. Then a graph P is called an *(x, y)-path* (or simply, a *path*) if P is a $\subseteq^s$-minimal graph in $\mathcal{G}_2^{1,1}$. Obviously, $x \neq y$. We call vertices x and y the *end-vertices of path P* and put $End(P) = \{x, y\}$.

Let $\mathcal{G}_2$ denote the set of graphs with all vertices having degree two. Then a graph C is called a *cycle* if C is a $\subseteq^s$-minimal graph in $\mathcal{G}_2$.

Let $\mathcal{G}_2^{1,3}$ denote the set of graphs having exactly one vertex, say x, of degree one, exactly one vertex, say y, of degree three, and all other vertices of degree two. Then a graph Q is called a *lollypop* if Q is a $\subseteq^s$-minimal graph in $\mathcal{G}_2^{1,3}$. We call vertex x the *end-vertex of lollypop Q*, put $End(Q) = \{x\}$, and also call Q an *x-lollypop*.



Figure 3: (x, y)-paths, x-lollypops, and x-cycles.

In a plain language an (x, y)-path is a graph $P = (V, E, \phi)$ with $V = \{x = v_1, \ldots, v_n = y\}$ and $E = \{e_1, \ldots, e_{n-1}\}$ and $\phi(e_i) = \{v_i, v_{i+1}\}$ for $i \in \{1, \ldots, n-1\}$. A graph G is a cycle if and only if G can be obtained from an (x, y)-path by adding a new edge e incident to x and y. A graph Q is a lollypop if and only if Q is obtained from disjoint cycle C (possibly, a loop) and a path P by identifying a vertex of C and an end-vertex of P.

Let $\mathcal{C}l(G)$ denote the set of subgraphs of G that are cycles. Let $\mathcal{C}t(G)$ denote the set of edge sets of elements in $\mathcal{C}l(G)$.

Given a graph G and edge e with the end vertices x and y, we say that *graph G' is obtained from by a subdivision of edge e* if $G' = (G \cup xPy) \setminus e$, where xPy is a path and $V(G) \cap V(P) = \{x, y\}$. A graph H is a *subdivision of graph G* if H is obtained from G by subdivisions of some of its edges.

A *k-skein*, $k \geq 3$, is a subdivision of a graph with two vertices and k parallel edges. A 3-skein is also called a Θ-*graph* (see Figure 4).

A *dumbbell* is a graph obtained from two disjoint cycles C, C', and a path xPx' disjoint from C and C' by identifying vertex x with a vertex in C and vertex x' with a vertex in C'

(see Figure 4).

A *butterfly* is a graph obtained from two disjoint cycles C, C' by identifying one vertex from C with one vertex in C' (see Figure 4).

Let $\mathcal{G}_2^{3,3}$ denote the set of graphs having exactly two vertices of degree three and all other vertices of degree two. Then, obviously, a $\subseteq^s$-minimal graph in $\mathcal{G}_2^{3,3}$ is either a Θ-graph or a dumbbell.

Let $\mathcal{G}_2^4$ denote the set of graphs having exactly one vertex of degree four and all other vertices of degree two. Then a $\subseteq^s$-minimal graph in $\mathcal{G}_2^4$ is a butterfly.

A graph G is *connected* if $|V(G)| \geq 2$ and for every two distinct vertices x and y in G there exists an (x, y)-path that is a subgraph of G.

A graph G is *2-connected* if $|V(G)| \geq 2$, G is connected and has no loops, and one of the following holds:
$(a1)$ if $|V(G)| = 2$, then $|E(G)| \geq 2$ (i.e. G has at least two parallel edges) and
$(a2)$ if $|V(G)| \geq 3$, then $G \setminus v$ is connected for every $v \in V(G)$.

A *multi-triangle* is either K_3 or a graph obtained from K_3 by replacing some edges by parallel edges.

A graph G is *3-connected* if $|V(G)| \geq 4$, G has no loops and no parallel edges, and $G \setminus v$ is 2-connected for every $v \in V(G)$.

A graph G is *multi-3-connected* if G is either 3-connected or can be obtained from a 3-connected graph by replacing some of its edges by parallel edges.

A graph G is *k-connected* for $k \geq 4$ if $|V(G)| \geq k + 1$, and $G \setminus v$ is $(k - 1)$-connected for every $v \in V(G)$.

A graph G is *of connectivity* k for $k \geq 1$ if G is k-connected but not $(k + 1)$-connected.

We call a graph G *cycle connected* if $|E(G)| \geq 2$ and for every two edges a and b in G there exists a cycle C in G such that $a, b \in E(C)$.

A *component of a graph* G is a $\subseteq^s$-maximal connected subgraph of G. Let $Cmp(G)$ denote the set of components of G and let $|Cmp(G)| = cmp(G)$.

A graph G is called *cacti-graph* if G has no isolated vertices, no leaves, and no cycle components. A connected cacti-graph is called a *cactus*. Let $\mathcal{G}_{\bowtie}$ denote the set of cacti-graphs and $\mathcal{CG}_{\bowtie}$ denote the set of connected graphs from $\mathcal{G}_{\bowtie}$, and so each member of $\mathcal{CG}_{\bowtie}$ is a cactus (for example, the graphs in Figures 8 and 9 and their subdivisions). Given a graph G, let $\mathcal{G}_{\bowtie}(G)$ denote the set of subgraphs of G that are members of $\mathcal{G}_{\bowtie}$. The class $\mathcal{G}_{\bowtie}$ of cacti-graphs will play a special role in our further discussion. Obviously, if a graph G is in $\mathcal{G}_{\bowtie}$, then every component of G is also in $\mathcal{G}_{\bowtie}$. Therefore every graph that is $\subseteq^s$-minimal in $\mathcal{G}_{\bowtie}$ is also $\subseteq^s$-minimal in $\mathcal{CG}_{\bowtie}$. In other words, a graph G is $\subseteq^s$-minimal in $\mathcal{G}_{\bowtie}$ if and only if G is $\subseteq^s$-minimal in $\mathcal{CG}_{\bowtie}$. Below (see Claim 3.2.4) we will show that G is $\subseteq^s$-minimal in $\mathcal{G}_{\bowtie}$ if and only if G is either a Θ-graph or a dumbbell or a butterfly.



3-skein ***dumbbell*** ***butterfly***

Figure 4: The structure of the minimal graphs in $\mathcal{G}_{\bowtie}$.

A subset X of vertices (of edges) in a graph G is called a *vertex cut* (respectively, an *edge cut*) of G if $G \setminus X$ has more components than G. A vertex cut (an edge cut) of G consisting of one element is called a *cut-vertex* (respectively, a *cut-egde* or a *bridge*) of G.

It is easy to prove the following claim.

Claim 2.3.1 *Let G be a graph and G' a graph obtained from G by deleting all isolated vertices. Then G is cycle connected if and only if G' is 2-connected.*

A *tree* is a connected graph without cycles. A *wheel* $W = (V, E, \phi)$ is a simple graph obtained from a cycle C with at least three vertices by adding a new vertex x (the *center of* W) and the set of edges $\{e_v : v \in V(C) \text{ and } \phi(e_v) = \{v, x\}\}$.

Claim 2.3.2 *Let G be a graph with $v(G) = v$ and $e(G) = e$. Suppose that G is 3-connected and has a vertex x such that $d(x, G) > e - v$. Then G is the wheel with center x, and so $d(x, G) = v - 1$ and $d(z, G) = 3$ for every $z \in V(G) \setminus x$.*

Proof Since $d(x, G) > e - v$, clearly $e(G \setminus x) < e - (e - v) = v$ and $v(G \setminus x) = v - 1$. Since G is 3-connected, $G - x$ is 2-connected. Therefore $e(G \setminus x) = v - 1$, $G \setminus x$ is a cycle, and vertex x is adjacent to every vertex of $G \setminus x$. Thus, G is the wheel with center x. $\square$

Using the arguments similar to those in the above proof it is also easy to prove the following claim.

Claim 2.3.3 *Let G be a graph. Suppose that G is 2-connected and has a vertex x such that $d(x, G) > e(G) - v(G) + 1$. Then every cycle of G contains x.*

From Claims 2.3.2 and 2.3.3 we have:

Claim 2.3.4 *Let G be a graph with $v(G) = v$ and $e(G) = e$. Suppose that G is 3-connected and not a complete graph on n vertices. Then one of the following holds:*

(c1) either every vertex of G has degree at most $e - v$ or

(c2) G has exactly one vertex x of degree $e - v + 1$ and every other vertex has degree at most $e - v$.

Graphs $G = (V, E, \phi)$ and $G' = (V', E', \phi')$ are *equal* if $V = V'$, $E = E'$, and $\phi = \phi'$.

An *isomorphism from* $G = (V, E, \phi)$ *to* $G' = (V', E', \phi')$ is a pair (ν, ε), where $\nu : V \to V'$ and $\varepsilon : E \to E'$ are bijections such that $\phi(e) = \{x, y\} \Leftrightarrow \phi'(\varepsilon(e)) = \{\nu(x), \nu(y)\}$. Graphs G and G' are *isomorphic* (denoted by $G \approx G'$) if there exists an isomorphism from G to G' (or, equivalently, an isomorphism from G' to G).

Let $G = (V, E, \phi)$ and $G' = (V', E', \phi')$ with a bijection $\varepsilon : E \to E'$. Then G and G' are called *strongly ε-isomorphic* if there exists a bijection $\nu : V \to V'$ such that (ν, ε) is an isomorphism from G to G'.

Without loss of generality, we may (and will) consider strongly isomorphic graphs G and G' with $E = E'$ (i.e. when $\varepsilon : E \to E'$ is the identity function). Namely, given two graphs $G = (V, E, \phi)$ and $G' = (V', E', \phi')$ with $E = E'$, a *strong isomorphism from* G to G' is a bijection $\nu : V \to V'$ such that $\phi(e) = \{x, y\} \Leftrightarrow \phi'(e) = \{\nu(x), \nu(y)\}$. Graphs $G = (V, E, \phi)$ and $G' = (V', E', \phi')$ with $E = E'$ are *strongly isomorphic* if there exists a strong isomorphism from G to G' (see Figures 5 and 6).



$$A \approx^s B \not\approx^s C$$

$$A \approx^s B \textit{ if } A \textit{ and } B \textit{ are strongly isomorphic}$$

Figure 5: Isomorphic and not strongly isomorphic graphs.

It is easy to prove the following claim.

Claim 2.3.5 Let G and G' be graphs and $E(G) = E(G')$. Then G and G' are strongly isomorphic if and only if $\mathcal{S}(G) = \mathcal{S}(G')$.

3 Some auxiliary notions and facts on graphs

3.1 On the Δ-function of a graph

Let $\mathcal{G}$ be the set of finite graphs and $G \in \mathcal{G}$. In this Section we will establish some properties of function $\Delta : \mathcal{G} \to \mathbb{Z}$ such that $\Delta(G) = |E(G)| - |V(G)|$. We will use these



Figure 6: Strongly isomorphic graphs.



Figure 7: The vertex stars of a graph

results on Δ in the study of so called k-circular matroids of a graph that will be defined later.

Instead of $\Delta(G)$ we will write simply ΔG. Let $X \subseteq E(G)$ and $G\langle X\rangle$ be the subgraph of G induced by X. Then $\Delta(G\langle X\rangle) = |X| - |V(G\langle X\rangle)|$.

We start with the following simple observation.

Claim 3.1.1 *Let G be a connected graph. If G has no cycle (i.e. G is a tree), then $\Delta G = -1$. If G has exactly one cycle, then $\Delta G = 0$. Graph G has at least two cycles if and only if $\Delta G \geq 1$.*

It is also easy to see the following:

Claim 3.1.2 *Let G be a graph. Then $\Delta G = \sum\{\Delta A : A \in Cmp(G)\}$.*

From Claims 3.1.1 and 3.1.2 we have:

Claim 3.1.3 *Let G be a graph and $t(G)$ the number of tree components of G. Then G has a component with at least two cycles if and only if $\Delta G + t(G) \geq 1$.*

Claim 3.1.4 *Let A be a connected subgraph of G and $\emptyset \neq X \subseteq E(A)$. Then $\Delta G\langle X\rangle \leq \Delta A$.*

Proof If $X = E(A)$, then our claim is obviously true. So we assume that $\emptyset \neq X \subset E(A)$. Let $S = V(A) \setminus V(G\langle X\rangle)$.

First, suppose that $S = \emptyset$. Since $|X| < |E(A)|$, we have: $\Delta G\langle X\rangle = |X| - |V(G\langle X\rangle)| < |E(A)| - |V(A)| = \Delta A$. Therefore our claim is true.

Finally, suppose that $S \neq \emptyset$. Since $X \neq \emptyset$ and A is connected, there are at least $|S|$ edges of $E(A) \setminus X$ that are incident to S, and so $|E(A) \setminus X| \geq |S|$. Then

$$\Delta A = |E(A)| - |V(A)| = |X| + |E(A) \setminus X| - (|V(G\langle X\rangle)| + |S|) =$$
$$\Delta G\langle X\rangle + |E(A) \setminus X| - |S| \geq \Delta G\langle X\rangle. \qquad \square$$

Claim 3.1.5 *Let F be a subgraph of G such that no component of F is a tree. Then $\Delta G\langle X\rangle \leq \Delta F$ for every subset X of $E(F)$.*

Proof (uses Claim 3.1.4)

Let $X \subseteq E(F)$ and $Cmp(F)$ be the set of components of F.
Let $\mathcal{R} = \{A \in Cmp(F) : E(A) \cap X \neq \emptyset\}$. Then $\Delta G\langle X\rangle = \sum_{A \in \mathcal{R}} \Delta G\langle E(A) \cap X\rangle$.
By Claim 3.1.4, $\Delta G\langle E(A) \cap X\rangle \leq \Delta A$ for every $A \in \mathcal{R}$.
Also, since every component of F is not a tree, $\Delta A \geq 0$ for every $A \in Cmp(F) \setminus \mathcal{R}$.
Thus,

$$\Delta G\langle X\rangle = \sum\{\Delta G\langle E(A) \cap X\rangle : A \in \mathcal{R}\} \leq \sum\{\Delta A : A \in \mathcal{R}\}$$
$$\leq \sum\{\Delta A : A \in Cmp(F)\} = \Delta F. \qquad \square$$

Now we will describe a discrete analog of the classical Intermediate Value Theorem on a continuos function.

A pair $D = (V, E)$ is called a *directed graph* (or simply, a *digraph*) if V is a non-empty finite set and $E \subseteq \{(x, y) : \{x, y\} \subseteq V \text{ and } x \neq y\}$. The elements of V and E are called the *vertices* and *arcs* of D, respectively. A vertex v in D is called *minimal* (*maximal*) if $(x, v) \notin E$ (respectively, $(v, x) \notin E$) for every $x \in V$.

Given $n, m \in \mathbb{Z}$ and $n \leq m$, let $[n, m] = \{x \in \mathbb{Z} : n \leq x \leq m\}$.

Theorem 3.1.6 DISCRETE INTERMEDIATE VALUE THEOREM

Suppose that digraph $D = (V, E)$ and function $f : V \to \mathbb{Z}$ satisfy the following conditions:

($d1$) *D has no directed cycles,*

($d2$) *D has exactly one minimal vertex p and exactly one maximal vertex b, and*

($d3$) *$f(p) \leq f(b)$ and $|f(x) - f(y)| \leq 1$ for every $(x, y) \in E$.*

Let $R = \{f(v) : v \in V\}$. Then $[f(p), f(b)] \subseteq R$.

Proof Suppose, to the contrary, that $[f(p), f(b)] \not\subseteq R$, i.e. the exists $r \in [f(p), f(b)] \setminus R$. Obviously, $f(p) < r < f(b)\}$. Let $V' = \{v \in V : f(v) < r\}$. Clearly, $p \in V'$ and $b \notin V'$. Therefore $V' \neq \emptyset$. Let $D' = D \setminus (V \setminus V')$, and so $V(D) = V'$. Since $b \notin V'$, digraph D' has a maximal vertex, say x. Since $x \neq b$, clearly x is not a maximal vertex in D. Therefore there exists $z \in V \setminus V'$ such that (x, z) is an arc in D. Since $z \in V \setminus V'$, we have: $f(z) \geq r$. If $f(z) > r$, then $f(z) - f(x) \geq 2$. However, $(x, z) \in E \Rightarrow |f(x) - f(z)| \leq 1$, a contradiction. Therefore $f(z) = r$, and so $r \in R$, a contradiction. $\square$

Claim 3.1.7 *Let $G = (V, E, \phi)$ be a graph. If $X \subseteq E$ and $e \in X$, then*
$|\Delta G\langle X\rangle - \Delta G\langle X \setminus\ e\rangle| \leq 1$, *i.e.* $\Delta G\langle X\rangle - \Delta G\langle X \setminus\ e\rangle \in \{-1, 0, 1\}$.

Proof The graph $G\langle X \setminus\ e\rangle$ has one fewer edge than $G\langle X\rangle$ and either zero or one or two fewer vertices. $\square$

From Theorem 3.1.6 and Claim 3.1.7 we have, in particular:

Claim 3.1.8 Let $G = (V, E, \phi)$ and $\Delta G\ = \delta \geq -1$. If at least one edge in E is not a loop, then for each $i \in \{-1, \ldots, \delta\}$ there exists $X \subseteq E$ such that $\Delta G\langle X\rangle = i$.

Proof (uses Theorem 3.1.6 and Claim 3.1.7)

Let $\mathcal{D} = (\mathcal{V}, \mathcal{E})$ be the digraph such that $\mathcal{V} = 2^E$ and if $X, Z \in \mathcal{V}$, then $(X, Z) \in \mathcal{E} \Leftrightarrow X \subset Z$ *and* $|Z \setminus X| = 1$. Obviously, digraph $\mathcal{D}$ has no directed cycles and has exacly one minimal vertex $p = \{\emptyset\}$ and exactly one maximal vertex $b = \{E\}$. Therefore digraph $\mathcal{D}$ satisfies conditions $(d1)$ and $(d2)$ of Claim 3.1.6.

Let function $f : \mathcal{V} \to \mathbb{Z}$ be defined as follows: $f(X) = \Delta G\langle X\rangle$ for $X \subseteq E$ and $X \neq \emptyset$, and so $f(E) = \delta$. Put $f(\emptyset) = -1$.

Obviously, for $e \in E$ we have: $f(\{e\}) = -1$ if e is not a loop and $f(\{e\}) = 0$ if e is a loop. Hence in both cases $|f(\{e\}) - f(\emptyset)| \leq 1$. Also if $X \subset E$ and $|X| \geq 2$, then by Claim 3.1.7, $|f(X) - f(X \setminus e)| \leq 1$. Thus, function $f : \mathcal{V} \to \mathbb{Z}$ satisfies condition $(d3)$ in Theorem 3.1.6. Therefore our claim follows from Theorem 3.1.6. $\square$

3.2 Splitter theorems for cacti-graphs

First we will remind the following known theorem which is very easy to prove.

Theorem 3.2.1 SPLITTER THEOREM FOR 2-CONNECTED GRAPHS
Let G be a 2-connected graph and G_0 a proper 2-connected subgraph of G. Then there exists a sequence $(P_1, \ldots, P_r)$ of paths in G and a sequence $(G_0, G_1, \ldots, G_r)$ of 2-connected subgraphs of G such that $G_r = G$, and $G_i = G_{i-1} \cup P_i$, where $V(G_{i-1}) \cap V(P_i) = End(P_i)$ for every $i \in \{1, \ldots, r\}$, and so $G_{i-1} \subset^s G_i$ and $\Delta G_i - \Delta G_{i-1} = 1$.

As in Section 2.3, $\mathcal{G}_{\bowtie}$ ($\mathcal{CG}_{\bowtie}$) is the set of graphs (respectively, connected graphs) with no isolated vertices, no leaves, and no cycle components, i.e. the set of cacti-graphs, (respectively, connected cacti-graphs). Notice that if graph G is 2-connected and it is not a cycle, then $G \in \mathcal{CG}_{\bowtie}$.

It turns out that similar Splitter Theorems are also true for the graph classes $\mathcal{CG}_{\bowtie}$ and $\mathcal{G}_{\bowtie}$. These theorems will be important for our further discussion.

Theorem 3.2.2 Splitter theorem for a cactus

Let G and G_0 be connected graphs. Suppose that $G \in \mathcal{CG}_{\bowtie}$ and G_0 is a subgraph of G such that either $G_0 \in \mathcal{CG}_{\bowtie}$ or G_0 is a cycle. Then there exist sequences $(P_1, \ldots, P_r)$ and $(G_0, G_1, \ldots, G_r)$ of subgraphs of G such that $G_r = G$, every $G_i \in \mathcal{CG}_{\bowtie}$, every P_i is either a path or a cycle or a lollipop (see Figure 3)*, and for every $i \in \{1, \ldots, r\}$, we have: $G_i = G_{i-1} \cup P_i$, where*

(**p**) *$V(G_{i-1}) \cap V(P_i) = End(P_i)$ if P_i is a a path,*

(**l**) *$V(G_{i-1}) \cap V(P_i) = End(P_i)$ if P_i is a lollipop, and*

(**c**) *$|V(G_{i-1}) \cap V(P_i)| = 1$ if P_i is a cycle,*

and so $G_{i-1} \subset^s G_i$ and $\Delta G_i - \Delta G_{i-1} = 1$.

We call the sequence $(G_0 \subset^s G_1 \subset^s \ldots \subset^s G_r = G)$ in the above theorem an *ear-assembly of a graph G in $\mathcal{G}_{\bowtie}$ starting from G_0* and each P_i an *ear of this ear-assembly.*

Proof of Theorem 3.2.2. We will first consider the case when $G_0 \in \mathcal{CG}_{\bowtie}$. The case when G_0 is a cycle will follow easily from the first case.

(**p1**) Suppose that $G_0 \in \mathcal{CG}_{\bowtie}$. Suppose, to the contrary, that our claim is not true. Our theorem is vacuously true if $G = G_0$. Suppose that G_k is a maximal subgraph of G such that our claim is true for G_k, and so there exist sequences $(P_1, \ldots, P_k)$ and $(G_0, \ldots, G_k)$ satisfying all the conditions of our theorem. Then $G_k \in \mathcal{CG}_{\bowtie}$ and G_k is a proper subgraph of G (i.e. $G_k \subset^s G$). We will get a contradiction by showing that there exist subgraphs P_{k+1} and G_{k+1} of G such that $G_k \cup P_{k+1} = G_{k+1}$, $G_{k+1} \in \mathcal{G}_{\bowtie}$, and pair (G_k, P_{k+1}) satisfies one of conditions (**p**),(**l**), (**c**).

Since G is connected and G_k is a proper subgraph of G, there is a path xPy in G such that $x \in V(G_k)$ and $V(P) \cap V(G_k) \subseteq \{x, y\}$. We can assume that xPy is a maximal path with the above mentioned properties.

First, suppose that $V(P) \cap V(G_k) = \{x, y\}$. Put $xPy = P_{k+1}$ and $G_{k+1} = G_k \cup P_{k+1}$. Then $G_{k+1} \in \mathcal{CG}_{\bowtie}$, $G_{k+1} \subseteq^s G$, and pair (G_k, P_{k+1}) satisfies condition (**p**).

Finally, suppose that $V(P) \cap V(G_k) = x$. Since $G \in \mathcal{CG}_{\bowtie}$, vertex y is not a leaf in G. Hence there exists edge $e = yz$ in $G \setminus (G_k \cup P)$. Since xPy is a maximal path with the above mentioned properties, vertex z is in P.

If $z \neq x$, then $P \cup e$ is a lollipop. Put $P_{k+1} = P \cup e$ and $G_{k+1} = G_k \cup P_{k+1}$. Then $G_{k+1} \in \mathcal{CG}_{\bowtie}$, $G_{k+1} \subseteq^s G$, and pair (G_k, P_{k+1}) satisfies condition (**l**).

If $z = x$, then $P \cup e$ is a cycle. Put $P_{k+1} = P \cup e$ and $G_{k+1} = G_k \cup P_{k+1}$. Then $G_{k+1} \in \mathcal{CG}_{\bowtie}$, $G_{k+1} \subseteq^s G$, and pair (G_k, P_{k+1}) satisfies condition (**c**).

(**p2**) Now suppose that G_0 is a cycle. The arguments similar to those in (**p1**) show that there exist subgraphs P_1 and G_1 of G such that $G_0 \cup P_1 = G_1$, $G_1 \in \mathcal{CG}_{\bowtie}$, and pair (G_0, P_1) satisfies one of conditions (**p**),(**l**), (**c**). Now put in (**p1**) $G_0 := G_1$. Then by (**p1**), our theorem is true for $G_0 := G_1$, i.e. there exists ear assembly $(G_1 \subset^s G_2 \subset^s \ldots \subset^s G_r = G)$. Then $(G_0 \subset^s G_1 \subset^s \ldots \subset^s G_r = G)$ is an ear assembly in $\mathcal{CG}_{\bowtie}$ of G starting from G_0. □

In order to formulate and prove a Splitter Theorem for $\mathcal{G}_{\bowtie}$, we need some preliminaries.

Definition 3.2.3 A graph D is a *bicycle* if D is a connected graph with no leaves and with $\Delta D = 1$.

Now we can describe the structure of graph-bicycles.

Claim 3.2.4 *A graph D is a bicycle if and only if D is either a Θ-graph or a dumbbell or a butterfly* (see Figure 4).

Proof (uses Theorem 3.2.2)
Obviously, a Θ-graph, a dumbbell, and a butterfly is a bicycle. It remains to prove that every bicycle D is either a Θ-graph or dumbbell or a butterfly. Since D is a bicycle, D is connected, has no leaves, and $\Delta D = 1$. Therefore $D \in \mathcal{CG}_{\bowtie}$ and $\Delta D = 1$, and so D has a cycle C and $\Delta C = 0$. Therefore our claim follows from Theorem 3.2.2, when G_0 is a cycle, $G_1 = D$, and $r = 1$. $\square$

It turns out that every graph G in $\mathcal{G}_{\bowtie}$ has bicycles and, moreover, graph G has some special bicycles.

Claim 3.2.5 *Suppose that G is a graph in $\mathcal{CG}_{\bowtie}$. Then for every two edges a, b in G there exists a bicycle in G containing a and b.*

Proof By Claim 3.2.4, it is sufficient to prove that G has either a Θ-graph or a dumbbell or a butterfly containing a and b.

(**p1**) Suppose that G has two cycles C, C' such that $a, b \in E(C \cup C')$. We can assume that $\{a, b\} \cap E(C) \neq \emptyset$.

Suppose first that $|V(C) \cap V(C')| \geq 2$. Then in C' there exists a path P such that $V(C) \cap V(P) = End(P)$ and $D = C \cup P$ contains both a and b. Then D is a Θ-graph in G containing a and b.

Now suppose that $|V(C) \cap V(C')| \leq 1$. Since G is connected, there exists a path xPx' in G (possibly, $x = x'$) such that $V(C) \cap V(P) = x$ and $V(C') \cap V(P) = x'$. If $|V(P)| = 1$, then $D = C \cup C'$ is a butterfly in G containing a and b. If $|V(P)| \geq 2$, then $D = C \cup P \cup C'$ is a dumbbell in G containing a and b.

(**p2**) Next suppose that G has a cycle C containing exactly one of a, b, say a, and b belongs to no cycle in G. Since G is connected, there exists a path xPx' such that $b \in E(P)$ and $V(C) \cap V(P) = x$. Since $G \in \mathcal{G}_{\bowtie}$, clearly G has no leaves, and therefore P belongs to a lollypop L in G such that $V(C) \cap V(L) = End(L)$. Since b belongs to no cycle in G, graph $C \cup L$ is a dumbbell containing both a and b.

(**p3**) Finally, suppose that neither a nor b belongs to a cycle in G. Since G is connected, there exists a path P in G containing a and b. Since $G \in \mathcal{G}_{\bowtie}$, clearly G has no leaves, and therefore P belongs to two lollypops L and L' in G. Then $L \cup P \cup L'$ is a dumbbell containing a and b. $\square$

Now we will formulate and prove a Splitter Theorem for graph class $\mathcal{G}_{\bowtie}$.

Theorem 3.2.6 SPLITTER THEOREM FOR A CACTI-GRAPH

Let $G, G_0 \in \mathcal{G}_{\bowtie}$. Suppose that G_0 is a subgraph of G. Then either $G = G_0$ or there exist sequences $(P_1, \ldots, P_r)$ and $(G_0, G_1, \ldots, G_r)$ of subgraphs of G such that $G_r = G$, every $G_i \in \mathcal{G}_{\bowtie}$, every P_i is either a path or a cycle or a lollipop (see Figure 3) *or a bicycle, and for every $i \in \{1, \ldots, r\}$, we have: $G_i = G_{i-1} \cup P_i$, where*

(**p**) *$V(G_{i-1}) \cap V(P_i) = End(P_i)$ if P_i is a a path,*

(**l**) *$V(G_{i-1}) \cap V(P_i) = End(P_i)$ if P_i is a lollipop,*

(**c**) *$|V(G_{i-1}) \cap V(P_i)| = 1$ if P_i is a cycle, and*

(**b**) *$V(G_{i-1}) \cap V(P_i) = \emptyset$ if P_i is a bicycle,*

and so $G_{i-1} \subset^s G_i$ and $\Delta G_i - \Delta G_{i-1} = 1$.

Proof (uses Theorem 3.2.2)

Suppose, to the contrary, that our claim is not true. Our theorem is vacuously true if $G = G_0$. Suppose that G_k is a maximal subgraph of G such that our claim is true for G_k, and so there exist sequences $(P_1, \ldots, P_k)$ and $(G_0, \ldots, G_k)$ satisfying all the conditions of our theorem. Then $G_k \in \mathcal{G}_{\bowtie}$ and G_k is a proper subgraph of G (i.e. $G_k \subset^s G$). We will get a contradiction by showing that there exist subgraphs P_{k+1} and G_{k+1} of G such that $G_k \cup P_{k+1} = G_{k+1}$, $G_{k+1} \in \mathcal{G}_{\bowtie}$, and pair (G_k, P_{k+1}) satisfies one of conditions (**p**),(**l**), (**c**), (**b**).

(**p1**) Suppose that G has a component A disjoint from G_k. Since $G \in \mathcal{G}_{\bowtie}$, also $A \in \mathcal{G}_{\bowtie}$. By Claim 3.2.5, A has a subgraph-bicycle B. Put $P_{k+1} = B$ and $G_{k+1} = G_k \cup P_{k+1}$. Then $G_{k+1} \in \mathcal{G}_{\bowtie}$, $G_{k+1} \subseteq^s G$, and pair (G_k, P_{k+1}) satisfies condition (**b**).

(**p2**) Now suppose that G has no component disjoint from G_k. Since G_k is a proper subgraph of G, there are components A of G and A_k of G_k such that A_k is a proper subgraph of A. Since $G, G_k \in \mathcal{G}_{\bowtie}$, also $A, A_k \in \mathcal{G}_{\bowtie}$. Therefore $A, A_k \in \mathcal{CG}_{\bowtie}$. Now our claim follows from Theorem 3.2.2 applied to A and A_k. □

We call the sequence $(G_0 \subset^s G_1 \subset^s \ldots \subset^s G_r = G)$ in the above theorem an *ear-assembly of a graph G in $\mathcal{G}_{\bowtie}$ starting from G_0* and each P_i an *ear* of this ear-assembly.

From Splitter Theorem 3.2.6 and Claim 3.2.5 we also have the following useful specification of this theorem.

Theorem 3.2.7 *Let $G \in \mathcal{G}_{\bowtie}$ and a, b edges in G.*

If a and b are in the same component of G, then either $G = G_0$ or there exists an ear-assembly $(G_0 \subset^s G_1 \subset^s \ldots \subset^s G_r = G)$ of graph G such that a and b are edges of G_i, $G_i \in \mathcal{G}_{\bowtie}$, and $\Delta G_i = i + 1$ for every $i \in \{0, \ldots, r\}$, and so, in particular, G_0 is a bicycle.

If a and b are in different components of G, then either $G = G_1$ or there exists an ear-assembly $(G_1 \subset^s G_2 \subset^s \ldots \subset^s G_r = G)$ of graph G such that a and b are edges of G_i, $G_i \in \mathcal{G}_{\bowtie}$, and $\Delta G_i = i + 1$ for every $i \in \{1, \ldots, r\}$.

Corollary 3.2.8 *Let $G \in \mathcal{G}_{\bowtie}$ and $a, b \in E(G)$.*

If a and b are in the same component of G, then for every $k \in \{1, \dots, \Delta G\}$ there exists a subgraph F of G such that $a, b \in E(F)$, $F \in \mathcal{G}_{\bowtie}$, and $\Delta F = k$.

If a and b are in different components of G, then for every $k \in \{2, \dots, \Delta G\}$ there exists a subgraph F of G such that $a, b \in E(F)$, $F \in \mathcal{G}_{\bowtie}$, and $\Delta F = k$.

Here is another characterization of bicycles.

Claim 3.2.9 A graph D is a bicycle if and only if D is a $\subseteq^s$-minimal graph in $\mathcal{G}_{\bowtie}$.

Proof (uses Theorem 3.2.6 and Claim 3.2.5)

(**p1**) Suppose that D is a bicycle. By Claim 3.2.4, D is either a Θ-graph or a dumbbell or a butterfly. It is easy to check that every such graph is in $\mathcal{G}_{\bowtie}$. Therefore $D \in \mathcal{G}_{\bowtie}$.

We prove that D is a $\subseteq^s$-minimal graph in $\mathcal{G}_{\bowtie}$. Suppose not. Then there exists graph D_0 in $\mathcal{G}_{\bowtie}$ such that D_0 is a proper subgraph of D. By Theorem 3.2.6, there exists an ear assembly $(D_0 \subset^s D_1 \subset^s \dots \subset^s D_r = D)$, where $\Delta D_i - \Delta D_{i-1} = 1$ for every $i \in \{1, \dots, r\}$. Therefore $\Delta D_0 < \Delta D$. Since $\Delta D = 1$, we have: $\Delta D_0 < 1$. Hence D_0 has a component with at most one cycle implying that $D_0 \not\in \mathcal{G}_{\bowtie}$, a contradiction. Therefore D is a $\subseteq^s$-minimal graph in $\mathcal{G}_{\bowtie}$.

(**p2**) Now suppose that D is a $\subseteq^s$-minimal graph in $\mathcal{G}_{\bowtie}$. Let A be a component of D. Since $D \in \mathcal{G}_{\bowtie}$, also $A \in \mathcal{G}_{\bowtie}$. By Claim 3.2.5, graph A contains a bicycle Q which is a subgraph of D. Since D is a $\subseteq^s$-minimal graph in $\mathcal{G}_{\bowtie}$, we have: $D = Q$, and so D is a bicycle. □

3.3 On the core and the kernel of a graph

In this Section we will give a definition and establish some properties of the *core* and the *kernel* of graphs.

For a graph A, let $\mathcal{R}_A = \{X \subseteq E(A) : X \neq \emptyset \textit{ and } \Delta A\langle X \rangle = \Delta A\}$.

Claim 3.3.1 *Let A be a connected graph with at least one cycle. Then $\mathcal{R}_A$ has the $\subseteq$-minimum element* (denoted by) $\mathcal{M}in(\mathcal{R}_A)$.

Proof (uses Claim 3.1.4 and 3.1.5)

Let $A_0 = A$ and define recursively A_s, for $s \geq 1$, to be the graph obtained from A_{s-1} by deleting all leaves in A_{s-1}. Consider $r \geq 0$ such that A_r has no leaves. Obviously, A_r is a connected graph, $E(A_r) \neq \emptyset$ and $\Delta A_r = \Delta A$. We claim that $E(A_r)$ is the $\subseteq$-minimum element in $\mathcal{R}_A$. Suppose, to the contrary, that $E(A_r)$ is not the $\subseteq$-minimum element in $\mathcal{R}_A$, i.e. there exists $X \in \mathcal{R}_A$ such that $E(A_r) \setminus X \neq \emptyset$. Since $X \in \mathcal{R}_A$, we have: $\emptyset \neq X \subseteq E(A)$ and $\Delta A\langle X \rangle = \Delta A$. Since $E(A_r) \setminus X \neq \emptyset$, there exists $e \in E(A_r) \setminus X$. If the graph $A \setminus e$ is connected, then by Claim 3.1.4, $\Delta A\langle X \rangle \leq \Delta(A \setminus e) < \Delta A$. If $A \setminus e$ is not connected, then $A \setminus e$ has two components. Since $e \in E(A_r)$, both components of $A \setminus e$ have cycles. Therefore, by Claim 3.1.5, $\Delta A\langle X \rangle \leq \Delta(A \setminus e) < \Delta A$. In both cases, $\Delta A\langle X \rangle < \Delta A$, a contradiction. □

Remark 3.3.2 If a graph A is a tree with at least two edges, then $\mathcal{R}_A$ has no $\subseteq$-minimum element.

Definition 3.3.3 THE KERNEL OF A GRAPH

If A is a connected graph with at least one cycle, then let $\lfloor A \rfloor$ denotes the subgraph of A induced by the edge subset $\mathcal{M}in(\mathcal{R}_A)$ defined in Claim 3.3.1, and so $\mathcal{M}in(\mathcal{R}_A) = E\lfloor A \rfloor$. If F is a non-connected graph, then put $\lfloor F \rfloor = \cup\{\lfloor A \rfloor : A \in Cmp(G) \text{ and } \Delta A \geq 0\}$.

Graph $\lfloor G \rfloor$ is called the *kernel of graph G*. If F is a forest, then the kernel of F is not defined.

Definition 3.3.4 THE CORE OF A GRAPH

Let G be a graph. Suppose that G has a component with at least two cycles. Put $[G] = \cup\{\lfloor A \rfloor : A \in Cmp(G) \text{ and } \Delta A \geq 1\}$. Graph $[G]$ is called *the core of graph G*. If every component of G has at most one cycle, then the core of G is not defined.

For a graph A, let $\mathcal{Q}_A = \{X \subseteq E(A) : A\langle X \rangle \textit{ has no leaves}\}$.

Claim 3.3.5 *Let A be a connected graph with at least one cycle. Then $\mathcal{Q}_A$ has the $\subseteq$-maximum element $\mathcal{M}ax(\mathcal{Q}_A)$. Moreover, $\mathcal{M}ax(\mathcal{Q}_A) = \mathcal{M}in(\mathcal{R}_A)$ and C is a subgraph of $\lfloor A \rfloor$ for every cycle C in A.*

Proof (uses Claim 3.3.1)

Let A_r be the graph defined in the proof of Claim 3.3.1. In that proof we have shown that $\lfloor A \rfloor = A_r$. It is clear that A_r has no leaves. Also, any subgraph of A containing $E(A_r)$ and at least one edge in $E(A) \setminus E(A_r)$ contains a leaf. Therefore, $E(A_r)$ is a $\subseteq$-maximal element in $\mathcal{Q}_A$. It is also clear that any set that is an element in $\mathcal{Q}_A$ is a subset of $E(A_r)$. Hence $E(A_r)$ is the $\subseteq$-maximum element in $\mathcal{Q}_A$. □

Let, as above,

$\mathcal{R}_G = \{X \subseteq E(G) : X \neq \emptyset \textit{ and } \Delta G\langle X \rangle = \Delta G\}$ and

$\mathcal{Q}_G = \{X \subseteq E(G) : G\langle X \rangle \textit{ has no leaves}\}$.

Claim 3.3.6 *Let G be a non-connected graph such that every component of G has at least two cycles. Then*

$$\mathcal{M}in(\mathcal{R}_G) = \cup\{\mathcal{M}in(\mathcal{R}_A) : A \in Cmp(G)\} =$$
$$\cup\{\mathcal{M}ax(\mathcal{Q}_A) : A \in Cmp(G)\} = \mathcal{M}ax(\mathcal{Q}_G) = E[G].$$

Proof (uses Claims 3.1.2 and 3.3.5)

The first equality follows easily from Claim 3.1.2. The second equality follows from Claim 3.3.5. The last equality follows from the fact that a graph has no leaf if and only if every component of the graph has no leaf. □

We recall that $\mathcal{G}_{\bowtie}$ is the set of graphs with no isolated vertices, no leaves, and no cycle components. Given a graph G, let $\mathcal{G}_{\bowtie}(G)$ denote the set of subgraphs of G that are members of $\mathcal{G}_{\bowtie}$.

From Claims 3.3.5 and 3.3.6 we have:

Theorem 3.3.7 Description of the core of a graph

Let G be a graph having a component with at least two cycles. The following are equivalent:

$(a1)$ *F is the core of G,*

$(a2)$ *F a unique $\subseteq^s$-maximum element in $\mathcal{G}_{\bowtie}(G)$, and*

$(a3)$ *if G' is the graph obtained from G by removing all tree components, then F is the only element in $\mathcal{G}_{\bowtie}(G)$ (or, equivalently, in $\mathcal{G}_{\bowtie}(G')$) such that $\Delta F = \Delta G'$.*

Remark 3.3.8 Let G be a graph. Then

$(c1)$ if G has a tree component, then $\mathcal{R}_G$ has no $\subseteq$-minimum element,

$(c2)$ if G is not connected, G has no tree component, and every component of G has exactly one cycle, then again $\mathcal{R}_G$ has no $\subseteq$-minimum element, and

$(c3)$ if G has no tree component but G has a component with exactly one cycle and also a component with at least two cycles, then $\mathcal{M}in(\mathcal{R}_G)$ exists and $\mathcal{M}in(\mathcal{R}_G) \subset \mathcal{M}ax(\mathcal{Q}_G)$.

Thus, Claim 3.3.6 provides a necessary and sufficient conditions for $\mathcal{M}in(\mathcal{R}_G) = \mathcal{M}ax(\mathcal{Q}_G)$.

Claim 3.3.9 *Let A be a connected graph with at least one cycle and $e \in E(A)$. Then*

$(c0)$ *$e \in E\lfloor A \rfloor$ if and only if both end vertices of e belong to $\lfloor A \rfloor$,*

$(c1)$ *if $e \notin E\lfloor A \rfloor$, then $A \setminus e$ has two components and exactly one of them is a tree and the other component contains $\lfloor A \rfloor$,*

$(c2)$ *if A has one cycle and $e \in E\lfloor A \rfloor$, then $A \setminus e$ is a tree,*

$(c3)$ *if A has at least two cycles and $e \in E\lfloor A \rfloor$, then every component of $A \setminus e$ contains a cycle, and*

$(c4)$ *if $A \setminus e$ has two components and $v \in V\lfloor A \rfloor$, then the component of $A \setminus e$ containing v is not a tree.*

Proof (uses Definition 3.3.3 and Claims 3.1.4 and 3.3.1)

(**p0**) We prove $(c0)$. Clearly, if $e \in E\lfloor A \rfloor$, then both end vertices of e belong to $\lfloor A \rfloor$. Now suppose that both end vertices of e belong to $\lfloor A \rfloor$ but $e \notin E\lfloor A \rfloor$. Then $\Delta(\lfloor A \rfloor \cup e) = \Delta\lfloor A \rfloor + 1 = \Delta A + 1$. Since $\lfloor A \rfloor \cup e$ is a subgraph of A, by Claim 3.1.4, $\Delta(\lfloor A \rfloor \cup e) \le \Delta A$, a contradiction.

(**p1**) We prove $(c1)$. First we claim that $A \setminus e$ is not connected. Suppose, to the contrary, $A \setminus e$ is connected. Then $\Delta(A \setminus e) = \Delta A - 1$. By Claim 3.1.4, $X \subseteq E(A \setminus e) \Rightarrow \Delta A\langle X \rangle \le \Delta A - 1$. Let $D = E\lfloor A \rfloor$. Then by Claim 3.3.1 and Definition 3.3.3, $\Delta A\langle D \rangle = \Delta A$. Therefore $E\lfloor A \rfloor = D \not\subseteq E(A \setminus e)$, and so $e \in E\lfloor A \rfloor$, a contradiction.

Now since A is connected and $A \setminus e$ is not connected, we have: $A \setminus e$ has two components A_1 and A_2, and clearly, one of them, say $A_2 = A \setminus A_1$, has a cycle. We claim that A_1 is a tree. Indeed, if not, then again $\Delta(A \setminus e) = \Delta A - 1$, and as above, $e \in E\lfloor A \rfloor$, a contradiction.

Since A_1 is a tree and e has one end-vertex in A_1 and the other end-vertex in A_2, we have: $\Delta A = \Delta(A \setminus A_1) = \Delta(A_2)$. Therefore, by Claim 3.3.1 and Definition 3.3.3, A_2 contains

$E\lfloor A\rfloor$.

(**p2**) We prove ($c2$). Since A has one cycle C, then $E\lfloor A\rfloor = E(C)$, and so $A \setminus e$ is a tree.

(**p3**) We prove ($c3$). If $A \setminus e$ is connected, then obviouly $A \setminus e$ contains a cycle. So we assume that $A \setminus e$ is not connected. Then $A \setminus e$ has two components, say A_1 and A_2 and e has one end-vertex in A_1 and the other end vertex in A_2, and so $e \notin E(A \setminus A_1)$. Clearly, at least one of A_1 and A_2, say $A_2 = A \setminus A_1$, contains a cycle.

Now, suppose to the contrary that A_1 has no cycle, i.e. A_1 is a tree.
Then $\Delta A = \Delta(A \setminus A_1) = \Delta(A_2)$. By Claim 3.3.1 and Definition 3.3.3, the core of A is an edge subset of A_2. Therefore $e \notin E\lfloor A\rfloor$, a contradiction.

(**p4**) Finally, we prove ($c4$).

First suppose that $e \notin E\lfloor A\rfloor$. Then by ($c1$), $A \setminus e$ has a component D containing $E\lfloor A\rfloor$. Since $\lfloor A\rfloor$ has at least one cycle, clearly D also has a cycle. Since vertex v is incident to $E\lfloor A\rfloor$, then v is in D.

Now suppose that $e \in E\lfloor A\rfloor$. Since $A \setminus e$ has two components, by ($c2$), A has at least two cycles. Then, by ($c3$), every component of $A \setminus e$ contains a cycle. □

From Claim 3.3.9 ($c1$) and ($c3$) we have:

Claim 3.3.10 *Let A be a connected graph with at least two cycles and $e \in E(A)$. Then the following are equivalent:*

($a1$) *$e \in E\lfloor A\rfloor = E[A]$ and*

($a2$) *every component of $A \setminus e$ has a cycle (or, equivalently, $A \setminus e$ has no tree component).*

From Claim 3.3.10 we have the following characterization of the edges in a graph belonging to the core.

Theorem 3.3.11 *Let G be a graph, $t(G)$ the number of tree components of G, and $Y(G) = \{a \in E(G) : t(G \setminus a) > t(G)\}$. Suppose that G has a component with at least two cycles.*
Then the following are equivalent:

($a1$) *$e \in E[G]$ and*

($a2$) *$e \in E(G) \setminus Y(G)$.*

Proof (uses Definition 3.3.4 and Claims 3.3.9 and 3.3.10)

If G is connected, then our claim follows from Claim 3.3.10. So we assume that G is not connected.

First, we prove ($a1$) ⇒ ($a2$). Suppose that ($a1$) holds, i.e. $e \in E[G]$. Then by Definition 3.3.4, $e \in E\lfloor A\rfloor$, where A is a component of G with at least two cycles and $\lfloor A\rfloor = [A]$. Then by Claim 3.3.10, $A \setminus e$ has no tree component, and so $t(G \setminus e) = t(G)$. Thus, $e \notin Y(G)$. Therefore ($a2$) holds.

Now we prove ($a2$) ⇒ ($a1$). Suppose that ($a2$) holds, i.e. $e \in E(G) \setminus Y(G)$. Then $t(G \setminus e) = t(G)$. Let A be the component of G containing e. Since $t(G \setminus e) = t(G)$, component A has at least two cycles. We claim that $e \in E\lfloor A\rfloor$. Suppose, not. Then by Claim 3.3.9 ($c1$), $A \setminus e$ has a tree component, say T. Then T is a component of $G \setminus e$. Clearly, T is not a component of G because T is a subgraph of the component A of G. Therefore $t(G \setminus e) > t(G)$, a contradiction. Thus, $e \in E\lfloor A\rfloor = E[A]$, and therefore $e \in E[G]$. Therefore ($a1$) holds. $\square$

4 On the k-circular matroid of a graph

In this Section we will introduce and study the properties of the so called *k-circular matroid $M_k(G)$ of a graph G*, where k is a non-negative integer. We will see, in particular, that (as before) $M_0(G)$ is the cycle matroid of graph G and $M_1(G)$ is the bicircular matroid of graph G.

The results of this Section will provide, in particular, a proper basis for our study of the problem $(WP)_k$ on describing the classes of graphs with the same k-circular matroids.

4.1 Circuits of the k-circular matroid of a graph

Definition 4.1.1 NON-DECREASING AND SUBMODULAR FUNCTIONS

Let E be a finite set. Consider function $f : 2^E \to \mathbb{Z}$.

Function f is called *non-decreasing* if $X \subseteq Y \subseteq E \Rightarrow f(X) \le f(Y)$.

Function f is called *submodular* if $X \subseteq Y \subseteq E \Rightarrow f(X \cup Y) + f(X \cap Y) \le f(X) + f(Y)$.

Theorem 4.1.2 (J. Edmonds and G.-C. Rota [4], see also [10])

Let E be a finite set and $f : 2^E \to \mathbb{Z}$ be a non-decreasing and submodular function. Then $\mathcal{C}(f) = \mathcal{M}in\{C \subseteq E : C \neq \emptyset$ and $|C| > f(C)\}$ is the collection of circuits of a matroid on E.

Notice that if $f(X) < 0$ for every $X \subseteq E$, then $\mathcal{C}(f) = \emptyset$, and therefore E is the only base of the matroid on E induced by function f.

The following is a simple strengthening of Theorem 4.1.2.

Claim 4.1.3 *Let E be a finite set and $f : 2^E \to \mathbb{Z}$ be a non-decreasing and submodular function. Then $\mathcal{C}(f) = \mathcal{M}in\ \{C \subseteq E : C \neq \emptyset$ and $|C| = f(C) + 1\}$.*

Proof Let $\mathcal{U} = \{C \subseteq E : C \neq \emptyset \text{ and } |C| > f(C)\}$ and $\mathcal{V} = \{C \subseteq E : C \neq \emptyset \text{ and } |C| = f(C) + 1\}$. It is clear that $\mathcal{V} \subseteq \mathcal{U}$. We want to show that $\mathcal{M}in\ \mathcal{U} = \mathcal{M}in\ \mathcal{V}$.

(**p1**) First we prove that $\mathcal{M}in\ \mathcal{U} \subseteq \mathcal{M}in\ \mathcal{V}$, i.e. that every minimal element of $\mathcal{U}$ is also a minimal element of $\mathcal{V}$. Let $X \in \mathcal{M}in\ \mathcal{U}$.

First we show that $X \in \mathcal{V}$. Since $X \in \mathcal{M}in\ \mathcal{U}$, clearly $X \in \mathcal{U}$, and therefore $X \neq \emptyset$. Let $e \in X$. By minimality of X in $\mathcal{U}$, $|X| - 1 = |X \setminus e| \le f(X \setminus e)$. Since f is a non-decreasing function, $f(X \setminus e) \le f(X)$. Since $X \in \mathcal{U}$, clearly $f(X) < |X|$. Now since in addition, f is an integer-valued function, we have: $f(X) \le |X| - 1$. Thus, $|X| - 1 = |X \setminus e| \le f(X \setminus e) \le f(X) \le |X| - 1$. It follows that $|X| = f(X) + 1$, and so $X \in \mathcal{V}$.

Now we will show that X is a minimal element of $\mathcal{V}$, i.e. that $X \in \mathcal{M}in\ \mathcal{V}$. Suppose, not. Then there exists $Y \subset X$ such that $Y \in \mathcal{V}$. Since $\mathcal{V} \subseteq \mathcal{U}$, we have: $Y \in \mathcal{V}$ and $Y \subset X$. Therefore $X \notin \mathcal{M}in\ \mathcal{U}$, a contradiction.

(**p2**) Now we will prove that $\mathcal{M}in\ \mathcal{U} \supseteq \mathcal{M}in\ \mathcal{V}$. Let $Y \in \mathcal{M}in\ \mathcal{V}$. Since $\mathcal{V} \subseteq \mathcal{U}$, clearly $Y \in \mathcal{U}$. Since $\mathcal{U}$ is a finite family, Y must contain a subset, say Z, which is a minimal element in $\mathcal{U}$. However, we have proved in (**p1**) that $\mathcal{M}in\ \mathcal{U} \subseteq \mathcal{M}in\ \mathcal{V}$, and therefore $Z \in \mathcal{M}in\ \mathcal{V}$. It follows that $Y = Z \in \mathcal{M}in\ \mathcal{U}$.

Thus, $\mathcal{M}in\ \mathcal{U} = \mathcal{M}in\ \mathcal{V}$. $\square$

Consider a function $f_k : 2^E \to \mathbb{Z}$ such that $f_k(X) = |V(G\langle X\rangle)| - 1 + k$ for every $X \in 2^E$. It is easy to see that the following is true.

Claim 4.1.4 ([4], see also [10])

Let $G = (V, E, \phi)$ be a graph and $k \geq 0$. Then f_k is a non-decreasing and submodular function.

From Claims 4.1.3 and 4.1.4 we have:

Claim 4.1.5 *Let $G = (V, E, \phi)$ be a graph and $k \geq 0$. Then*

$\mathcal{C}_k(G) = \mathcal{M}in\ \{C \subseteq E : C \neq \emptyset$ *and* $|C| = |V(G\langle C\rangle)| + k\}$

is the collection of circuits of a matroid on E.

Definition 4.1.6 THE k-CIRCULAR MATROID OF A GRAPH

We call the matroid described in Claim 4.1.5 by the set $\mathcal{C}_k(G)$ of its circuits *the k-circular matroid of G* and denote it $M_k(G)$. Let $\mathcal{D}_k(G)$, $\mathcal{I}_k(G)$, $\mathcal{B}_k(G)$, and $\mathcal{C}^*_k(G)$ denote the families of dependent sets, independent sets, bases, and cocircuits of $M_k(G)$, respectively.

It is easy to see that (as above) $M_0(G)$ is the cycle matroid of a graph G. As we will see below, matroid $M_1(G)$ is the bicircular matroid $B(G)$.

From Claim 4.1.5 we have:

Claim 4.1.7 *Let $G = (V, E, \phi)$ be a graph and $k \geq 0$. Then*

$\mathcal{C}_k(G) = \mathcal{M}in\ \{C \subseteq E : C \neq \emptyset$ *and* $\Delta G\langle C\rangle = k\}$.

As in Section 1, $\mathcal{G}_{\bowtie}$ is the set of graphs with no isolated vertices, no leaves, and no cycle components, i.e. the set of cacti-graphs.

Theorem 4.1.8 STRUCTURE OF GRAPH INDUCED BY A CIRCUIT OF $M_k(G)$ IN G

Let $G = (V, E, \phi)$ be a graph. Let $C \subseteq E$ and $k \geq 1$. Then the following are equivalent:

(c1) $C \in \mathcal{C}_k(G)$ and

(c2) $\Delta G\langle C\rangle = k$ and $G\langle C\rangle \in \mathcal{G}_{\bowtie}$, i.e. $G\langle C\rangle$ has no isolated vertices, no leaves, and every component of $G\langle C\rangle$ has at least two cycles.

Proof (uses Claim 4.1.7)

Obviously, a connected graph A has at least two cycles if and only if $\Delta A \geq 1$.

(**p1**) First we prove $(c1) \Rightarrow (c2)$. By Claim 4.1.7, $\Delta G\langle C\rangle = k$. Also, if e is a pendant edge of $G\langle C\rangle$, then clearly, $\Delta G\langle C\rangle - \Delta G\langle C \setminus e\rangle \in \{-1, 0\}$. Therefore $\Delta G\langle C \setminus e\rangle \geq \Delta G\langle C\rangle$, contradicting minimality of C. Hence $G\langle C\rangle$ contains no leaves.

Let $\mathcal{A}$ be the set of components of $G\langle C\rangle$. Clearly, $E(A) \neq \emptyset$ for every $A \in \mathcal{A}$, $C = \bigcup_{A\in\mathcal{A}} E(A)$, and $V(G\langle C\rangle) = \bigcup_{A\in\mathcal{A}} V(A)$. Then

$$\begin{aligned}\textstyle\sum_{A\in\mathcal{A}} \Delta A &= \textstyle\sum_{A\in\mathcal{A}}(|E(A)| - |V(A)|) = \sum_{A\in\mathcal{A}} |E(A)| - \sum_{A\in\mathcal{A}} |V(A)| \\ &= |C| - |V(G\langle C\rangle)| = \Delta G\langle C\rangle = k \geq 1.\end{aligned}$$

Therefore, $\sum_{A\in\mathcal{A}} \Delta A = \Delta G\langle C\rangle \geq 1$.

If $|\mathcal{A}| = 1$, then $\Delta A \geq 1$, where A is the unique component of $G[C]$. Now suppose that $|\mathcal{A}| \geq 2$ and let $A' \in \mathcal{A}$. Then the set of components of $G\langle C \setminus E(A')\rangle$ is $\mathcal{A}' = \mathcal{A} \setminus \{A'\}$. Therefore

$$\Delta G\langle C \setminus E(A')\rangle = \textstyle\sum_{A\in\mathcal{A}'} \Delta A = \sum_{A\in\mathcal{A}} \Delta A - \Delta A' = \Delta G\langle C\rangle - \Delta A'.$$

Thus, if $\Delta A' < 1$, then $\Delta G\langle C \setminus E(A')\rangle \geq \Delta G\langle C\rangle = k$, contradicting minimality of C. Hence every component A of $G\langle C\rangle$ has at least two cycles.

(**p2**) Finally, we prove $(c1) \Leftarrow (c2)$. Consider $C \subseteq E$ satisfying $(c2)$. Then $\Delta G\langle C\rangle = k$, where $k \geq 1$. Therefore $C \neq \emptyset$. Hence C is an element of set $\mathcal{V} = \{Z \subseteq E : Z \neq \emptyset \textit{ and } \Delta G\langle Z\rangle = k\}$. It remains to show that C is a minimal element in $\mathcal{V}$.

We first prove the following

Claim. *If A is a component of $G\langle C\rangle$ and $X \subset E(A)$, then $\Delta G\langle X\rangle < \Delta A$.*

Proof. If $X = \emptyset$, then $\Delta G\langle X\rangle = 0 < 1 \leq \Delta A$. So we assume that $X \neq \emptyset$.

Let $S = V(A) \setminus V(G\langle X\rangle)$. If $S = \emptyset$, then $V(G\langle X\rangle) = V(A)$ and since $|X| < |E(A)|$, we have: $\Delta G\langle X\rangle = |X| - |V(G\langle X\rangle)| < |E(A)| - |V(A)| = \Delta A$. So we assume that $S \neq \emptyset$. Since $X \neq \emptyset$ and A is connected and has no leaves, there are at least $|S|+1$ edges of $E(A)\setminus X$ that are incident to vertices in S, and so $|E(A) \setminus X| > |S|$. Then

$$\begin{aligned}\Delta G\langle E(A)\rangle &= E(A) - V(A) = |X| + |E(A) \setminus X| - (|V(G\langle X\rangle)| + |S|) \\ &= \Delta G\langle X\rangle + |E(A) \setminus X| - |S| \geq \Delta G\langle X\rangle + 1 > \Delta G\langle X\rangle. \qquad \diamondsuit\end{aligned}$$

Now we prove that C is a minimal element in $\mathcal{V}$, namely, that $Z \subset C$ *and* $Z \neq \emptyset \Rightarrow Z \notin \mathcal{V}$, i.e. $\Delta_G(Z) < k$.

Let $\mathcal{A}$ be the set of components of $G\langle C\rangle$. Put $Z_A = Z \cap E(A)$ for $A \in \mathcal{A}$. Then

$$\Delta G\langle Z\rangle = |Z| - |V(G\langle Z\rangle)| = \textstyle\sum_{A\in\mathcal{A}} |Z \cap E(A)| - \sum_{A\in\mathcal{A}} |V(G\langle Z\rangle) \cap V(A)| =$$
$$\textstyle\sum_{A\in\mathcal{A}}(|Z \cap\ E(A)| - |V(G\langle Z\rangle) \cap\ V(A)|) = \sum_{A\in\mathcal{A}} \Delta G\langle Z_A\rangle.$$

Since $Z \subset Y$, we have: $Z_A \subset A$ for some $A \in \mathcal{A}$. Now by the above Claim,

$$\Delta G\langle Z\rangle = \textstyle\sum_{A\in\mathcal{A}} \Delta G\langle Z_A\rangle < \ \sum_{A\in\mathcal{A}} \Delta A = \Delta G\langle C\rangle = k. \qquad \square$$

We remind that a graph R is a bicycle if and only if R is a connected graph with no leaves and $\Delta R = 1$ (see Figure 8).

From the definition of a bicycle, Claim 3.2.9, and Theorem 4.1.8 we have:

Claim 4.1.9 *$C \in \mathcal{C}_1(G)$ if and only if $G\langle C\rangle$ is a bicycle* (see Figure 8).

Remark 4.1.10 There is an essential difference between bicircular matroid $M_1(G)$ and k-circular matroid $M_k(G)$ for $k \geq 2$. Namely, every circuit of $M_1(G)$ induces a connected subgraph in G. Therefore if $M_1(G)$ is a connected matroid, then G is a connected graph. On the other hand, if $k \geq 2$, then $M_k(G)$ may have a circuit that induces a non-connected subgraph in G. Therefore for $k \geq 2$ matroid $M_k(G)$ may be connected although graph G is not connected.

A circuit C of $M_k(G)$ is called *graph-connected* if the graph induced by C in G is connected.

Matroid $M_2(G)$ is called a *tricircular matroid*. The connected circuits of tricircular matroids are subdivisions of graphs in Figure 9. The non-connected circuits of tricircular matroids are pairs of disjoint subdivisions of graphs from $\{S, D, B\}$ in Figure 8.



Figure 8: Bicycles (up to subdivisions).



The graph structures of the connected circuits of tricircular matroids

Figure 9: Connected tricycles (up to subdivisions).

Theorem 4.1.11 Recursive description of a circuit of $M_k(G)$

Let $G = (V, E, \phi)$ *be a graph. Let* $C \subseteq E$ *and* $k \geq 1$. *If* $k = 1$, *then* $C \in \mathcal{C}_1(G)$ *if and only if* $G\langle C\rangle$ *is a bicycle. If* $k \geq 2$, *then the following are equivalent:*

$(c1)$ $C \in \mathcal{C}_k(G)$ *and*

$(c2)$ *there exists* $C' \in \mathcal{C}_{k-1}(G)$ *such that* $G\langle C\rangle = G\langle C'\rangle \cup P$, *where* P *is either a path or a cycle or a lollipop* (see Figure 3) *or a bicycle and*

$(\mathbf{l}, \mathbf{p})$ $V(G\langle C'\rangle) \cap V(P) = End(P)$ *if* P *is a lollipop or a path,*

$(\mathbf{c})$ $|V(G\langle C'\rangle) \cap V(P)| = 1$ *if* P *is a cycle, and*

$(\mathbf{b})$ $V(G\langle C'\rangle) \cap V(P) = \emptyset$ *if* P *is a bicycle.*

Proof (uses Theorems 3.2.6 and 4.1.8 and Claim 3.2.7)

If $k = 1$, then by Claim 4.1.9, $C \in \mathcal{C}_1(G)$ if and only if $G\langle C\rangle$ is a bicycle. So we assume that $k \geq 2$.

First, we prove $(c1) \Rightarrow (c2)$. Let $C \in \mathcal{C}_k(G)$. Then by Theorem 4.1.8, $\Delta G\langle C\rangle = k$ and $G\langle C\rangle \in \mathcal{G}_{\bowtie}$. By Claim 3.2.7, graph $G\langle C\rangle$ contains a bicycle D. Therefore by Theorem 3.2.6, there exists a subgraph $G_{k-1} = G\langle C'\rangle$ such that $C' \in \mathcal{C}_{k-1}(G)$ and $(c2)$ holds.

Now we prove $(c2) \Rightarrow (c1)$. Since $(c2)$ holds, $G\langle C\rangle \in \mathcal{G}_{\bowtie}$ and $\Delta G\langle C\rangle = \Delta G\langle C'\rangle + 1 = k$. Therefore by Theorem 4.1.8, $C \in \mathcal{C}_k(G)$, and so $(c1)$ holds. □

Definition 4.1.12 Given two graphs G and G', we say that G and G' are M_k-*equal* and write $G \sim^k G'$ if $M_k(G) = M_k(G')$. Clearly, relation $\sim^k$ on the set of finite graphs is an equivalence relation. The equivalence classes of this relation are called M_k-*equivalence classes.*

Using Theorems 3.2.6 and 4.1.8 it is not difficult to prove the following claim about the relation between M_k- and M_s-equivalence classes of graphs for $k > s \geq 1$.

Claim 4.1.13 *Let* $k > s \geq 1$. *If* $\mathcal{F}$ *be an* M_k*-equivalence class of graphs, then* $\mathcal{F}$ *is the union of some* M_s*-equivalence classes of graphs. In other words, if* $M_s(G) = M_s(G')$, *then* $M_k(G) = M_k(G')$.

Claim 4.1.14 *Let* G *be a graph and* $k \geq 1$. *Then* $M_k(G)$ *has no loops.*

Proof (uses Theorem 4.1.8)

By Theorem 4.1.8, if C is a circuit of $M_k(G)$, then $G\langle C\rangle$ has at least two cycles. Therefore $|E(C)| \geq 2$. □

Claim 4.1.15 *Let* $G = (V, E, \phi)$ *and* $k \geq 1$. *If* $X \subseteq E$ *and* $\Delta G\langle X\rangle \geq k$, *then* $X \in \mathcal{D}_k(G)$.

Proof By Claim 3.1.8, the set $\mathcal{P} = \{Z \subseteq X : \Delta G\langle Z\rangle = k\}$ is non-empty. A minimal element of $\mathcal{P}$ exists and belongs to $\mathcal{C}_k(G)$, implying that $X \in \mathcal{D}_k(G)$. □

4.2 Non-trivial k-circular matroids

We remind that a matroid $M = (E, \mathcal{I})$ is non-trivial if E is not a base and not a cobase of M.

In this Subsection we give a criterion for a k-circular matroid of a graph to be non-trivial and, on the other hand, describe all graph representations of a trivial k-circular matroid of a graph.

Theorem 4.2.1 A criterion for matroid $M_k(G)$ to be non-trivial

Let G be a graph, $F(G)$ the union of all tree components of G, and $k \geq 1$. Then the following are equivalent:

$(a1)$ $\mathcal{C}_k(G) \neq \emptyset$,

$(a2)$ $k \leq \Delta G + cmp(F(G))$, *and*

$(a3)$ $M_k(G)$ *is a non-trivial matroid.*

Proof (uses Theorem 4.1.8 and Claims 3.1.5 and 4.1.15)

Let $G' = G \setminus F(G)$. Clearly, $\Delta G = \Delta G' + \Delta F(G) = \Delta G' - cmp(F(G))$. Therefore $\Delta G' = \Delta G + cmp(F(G))$.

First we prove $(a1) \Rightarrow (a2)$. Suppose that $(a1)$ is true, i.e. $\mathcal{C}_k(G) \neq \emptyset$. Let $C \in \mathcal{C}_k(G)$. Then by Theorem 4.1.8, $\Delta G\langle C\rangle \geq k$ and every component of $G\langle C\rangle$ has at least two cycles. Since every vertex of $F(G)$ belongs to a tree components of G, graphs $G\langle C\rangle$ and $F(G)$ have no vertex in common. Since $G' = G \setminus F(G)$, graph $G\langle C\rangle$ is a subgraph of G'. Since G' has no tree components, by Claim 3.1.5, $k \leq \Delta G\langle C\rangle \leq \Delta G' = \Delta G + cmp(F(G))$.

Next we prove $(a2) \Rightarrow (a1)$. Suppose that $(a1)$ is true, i.e. $k \leq \Delta G + cmp(F(G))$. Since $\Delta G' = \Delta G + cmp(F(G))$, we have $\Delta G' \geq k$. Then by Claim 4.1.15, $E(G') \in \mathcal{D}_k(G)$ and therefore $\mathcal{C}_k(G) \neq \emptyset$.

Now we prove $(a1) \Leftrightarrow (a3)$. By definition of a non-trivial matroid, $(a3) \Rightarrow (a1)$. By Claim 4.1.14, $M_k(G)$ has no loops, and therefore $E(G)$ is not a cobase of $M_k(G)$. By $(a1)$, $E(G)$ is not a base of $M_k(G)$. Thus, $M_k(G)$ is a non-trivial matroid, and therefore $(a1) \Rightarrow (a3)$. $\square$

From Theorem 4.1.8 and Claim 4.2.1 we have:

Theorem 4.2.2 *Let G be a graph. Then $M_k(G)$ is a trivial matroid if and only if G has no subgraph F such that $F \in \mathcal{G}_{\bowtie}$ and $\Delta F = k$.*

Remark 4.2.3 The above Theorem 4.2.2 provides for a given $k \geq 1$ a complete description of all graphs G that have the same matroid $M_k(G)$ in case when $M_k(G)$ is a trivial matroid. In other words, the above Theorem 4.2.2 describes all graph representations of the k-circular trivial matroid.

For this reason from now on we will consider the situations when $M_k(G)$ is a non-trivial matroid.

4.3 Coloops of non-trivial k-circular matroids

Now we are going to give a graph characterization of the set of coloops for a non-trivial matroid $M_k(G)$.

We know that an element e in a matroid M is a coloop of M if and only if e belongs to no circuit of M.

Obviously, from Claim 4.2.1 we have:

Claim 4.3.1 *$M_k(G)$ is a non-trivial matroid if and only there exists an edge in G which is not a coloop of $M_k(G)$, or, equivalently, $M_k(G)$ is a trivial matroid if and only if every edge of G is a coloop of $M_k(G)$.*

Claim 4.3.2 *Let G be a graph and $R = R(G)$ the union of components of G having at most one cycle. Then every edge e of R is a coloop of matroid $M_k(G)$ for every positive integer k, and so $E(R)$ is a subset of every base of $M_k(G)$.*

Proof (uses Claim 4.1.8)

Obviously, an element e of a matroid M is a coloop if an only if belongs to no circuit of M. Let $e \in E(R)$, i.e. e belongs to a component of G, say T, having at most one cycle.

It is sufficient to prove that e belongs to no circuit of $M_k(G)$. Suppose, to the contrary, that there exists a circuit C in $M_k(G)$ containing e. Let A be the component of $G\langle C\rangle$ containing e. Then by Claim 4.1.8, A has at least two cycles D_1 and D_2. Now since $e \in E(A) \cap E(T)$, clearly A is a subgraph the component T, and so cycles D_1 and D_2 are subgraphs of T. □

Notice that if $M_k(G)$ is a non-trivial matroid for some $k \geq 1$, then by Claim 4.2.1, there exists $C \in \mathcal{C}_k(G)$. By Theorem 4.1.8, every component of $G\langle C\rangle$ has at least two cycles. Therefore graph G has a component with at least two cycles, and so the core $[G]$ of graph G (see Definition 3.3.4) is defined.

Let $L^*_k(G)$ denote the set of coloops of $M_k(G)$.

Claim 4.3.3 *Let $k \geq 1$ and $E = E(G)$. Suppose that $M_k(G)$ is a non-trivial matroid. Then $L^*_k(G) = E \setminus E[G]$.*

Proof (uses Theorem 3.2.6, 3.3.7, and 4.1.8, Corollary 3.2.8, and Claims 3.3.5, 3.3.9, and 4.3.2)

(**p1**) First we prove that $E \setminus E[G] \subseteq L^*_k(G)$. Let $e \in E \setminus E[G]$ and A be the component of G containing e. If A has at most one cycle, then by Claim 4.3.2, $e \in L^*_k(G)$. So we assume that A has at least two cycles. Since $e \in E \setminus E[G]$, clearly $e \notin E[A] = E\lfloor A\rfloor$. Then, by Claim 3.3.9 ($c1$), graph $A \setminus e$ has a tree component. We claim that e belongs to no circuit of $M_k(G)$. Suppose, to the contrary, that there exists a circuit C of $M_k(G)$ containing e. Let D be the component of $G\langle C\rangle$ containing e. By Theorem 4.1.8, D has no leaf. Then by Claim 3.3.5, $e \in E(D) \subseteq E\lfloor A\rfloor$, a contradiction. Therefore e belongs to no circuit of $M_k(G)$, and so $e \in L^*_k(G)$. Hence, $E \setminus E[G] \subseteq L^*_k(G)$.

(**p2**) Now we prove that $L_k^*(G) \subseteq E \setminus E[G]$. Suppose, to the contrary, that there exists $e \in L_k^*(G)$ such that $e \notin E \setminus E[G]$, i.e. $e \in E[G]$. We will show that there exists $C \in \mathcal{C}_k(G)$ containing e, contradicting $e \in L_k^*(G)$.

Since $M_k(G)$ is a non-trivial matroid, there exists $D \in \mathcal{C}_k(G)$. By Theorem 4.1.8, $\Delta G\langle D\rangle = k \geq 1$ and $G\langle D\rangle \in \mathcal{G}_{\bowtie}$. Therefore G has a component with at least two cycles. Then by Theorem 3.3.7, $[G] \in \mathcal{G}_{\bowtie}$ and $G\langle D\rangle \subseteq^s [G]$. Also, by Theorem 3.2.6, $\Delta[G] \geq \Delta G\langle D\rangle = k$. Now by Corollary 3.2.8, there exists a subgraph F of $[G]$ such that $e \in E(F)$, $F \in \mathcal{G}_{\bowtie}$, and $\Delta(F) = k$. Put $E(F) = C$. Then $e \in C$ and by Theorem 4.1.8, $C \in \mathcal{C}_k(G)$. ☐

Given a graph G, let $t(G)$ be the number of tree components of G, and $Y(G) = \{e \in E(G) : t(G \setminus e) > t(G)\}$.

From Claims 3.3.11 and 4.3.3 we have the following more detailed characterization of the set of coloops of a non-trivial matroid $M_k(G)$.

Theorem 4.3.4 GRAPH DESCRIPTION OF COLOOPS OF $M_k(G)$

Let G be a graph and $M_k(G)$ a non-trivial matroid. Then $L_k^*(G) = E \setminus E[G] = Y(G)$.

4.4 Connected k-circular matroids

We recall that a matroid $M = (E, \mathcal{I})$ is connected if $|E| \geq 2$ and for every $a, b \in E$ there exists $C \in \mathcal{C}(M)$ such that $a, b \in C$.

In this Subsection we present a graph criterion for a k-circular matroid to be connected.

We start with the following claims.

Claim 4.4.1 *Let G be a graph for which $M_k(G)$ is a non-trivial matroid. Then $\mathcal{C}_k(G) = \mathcal{C}_k[G]$.*

Proof (uses Claims 2.2.3 and 4.3.3)

Obviously, $\mathcal{C}_k[G] \subseteq \mathcal{C}_k(G)$. We prove $\mathcal{C}_k(G) \subseteq \mathcal{C}_k[G]$. Let $C \in \mathcal{C}_k(G)$. Then by Claim 2.2.3, $C \cap L_k^*(G) = \emptyset$. By Claim 4.3.3, $C \cap (E \setminus E[G]) = \emptyset$. Therefore $C \subseteq E[G]$. ☐

From Claim 4.4.1 we have:

Claim 4.4.2 *Let G and G' be graphs and $k \geq 1$. Suppose that $M_k(G)$ is a non-trivial matroid. Then $M_k(G) = M_k(G')$ if and only if $E(G) = E(G')$ and $M_k[G] = M_k[G']$.*

Claim 4.4.3 *If $M_k(G)$ is a non-trivial matroid, then the core $[G]$ of G is defined.*

Proof (uses Theorems 3.2.7 and 4.1.8)

Since $M_k(G)$ is non-trivial, there exists $C \in \mathcal{C}_k(G)$. By Theorem 4.1.8, $G\langle C\rangle \in \mathcal{G}_{\bowtie}$. By Theorem 3.2.7, $G\langle C\rangle$ contains a bicycle, and so G has a component with at least two cycles. Therefore the core $[G]$ of G is defined. ☐

Remark 4.4.4 Claim 4.4.2 and Theorem 3.3.7 allows us to reduce our problem $(WP)_k$ of describing the classes of graphs with the same non-trivial k-circular matroid to the corresponding problem for the graphs in $\mathcal{G}_{\bowtie}$, i.e. for the class of cacti-graphs.

For this reasons from now on we will consider the situations when G is a cacti-graph.

Let $L_k(G)$ and $L_k^*(G)$ denote the set of loops and coloops of $M_k(G)$, respectively. By Claim 4.1.14, $L_k(G) = \emptyset$ and by Claim 4.3.3, $L_k^*(G) = E \setminus E[G]$ for any non-trivial $M_k(G)$. Therefore $E \setminus (L_k(G) \cup L_k^*(G)) = E \setminus L_k^*(G) = E[G]$. In particular, for cacti-graphs $L_k(G) \cup L_k^*(G)) = \emptyset$.

In what follows we give a characterization of a non-trivial matroid $M_k(G)$ for a cacti-graph G.

Theorem 4.4.5 GRAPH DESCRIPTION OF CONNECTED MATROID $M_k(G)$

Let $k \geq 2$. Then the following are equivalent:

$(a1)$ *$M_k(G)$ is a non-trivial matroid and $G \in \mathcal{G}_{\bowtie}$ and*

$(a2)$ *$M_k(G)$ is a connected matroid.*

Proof (uses Theorems 3.2.6, 3.3.7, and 4.1.8, Corollary 3.2.8, and Claims 4.3.3 and 4.4.3)

(**p1**) First we prove $(a1) \Rightarrow (a2)$.

Let $a, b \in E(G)$. Since $M_k(G)$ is a non-trivial matroid, there exists $D \in \mathcal{C}_k(G)$. By Theorem 4.1.8, $\Delta G\langle D\rangle = k$ and $G\langle D\rangle \in \mathcal{G}_{\bowtie}$. Since $G \in \mathcal{G}_{\bowtie}$ and $G\langle D\rangle \subseteq^s G$, by Theorem 3.2.6, $\Delta G \geq \Delta G\langle D\rangle = k$. By Corollary 3.2.8, there exists a subgraph F of G such that $a, b \in E(F)$, $F \in \mathcal{G}_{\bowtie}$, and $\Delta(F) = k$. Put $E(F) = C$. Then $a, b \in C$ and by Theorem 4.1.8, $C \in \mathcal{C}_k(G)$. Therefore $M_k(G)$ is a connected matroid.

(**p2**) Finally, we prove $(a2) \Rightarrow (a1)$. Since $M_k(G)$ is a connected matroid, we have: $M_k(G)$ has a k-circuit C, and therefore $M_k(G)$ is non-trivial. By Claim 4.4.3, the core $[G]$ of G is defined. Since $M_k(G)$ is a connected matroid, $L_k^*(G) = \emptyset$. Then by Claim 4.3.3, $E = E[G]$, and so $G = [G]$. Now by Theorem 3.3.7, $G \in \mathcal{G}_{\bowtie}$. $\square$

Theorem 4.4.6 GRAPH DESCRIPTION OF THE $\sim$- EQUIVALENCE CLASSES IN $M_1(G)$

Let G be a graph. Then the following are equivalent:

$(a1)$*$M_1(G)$ is a non-trivial matroid and $G \in \mathcal{G}_{\bowtie}$ and*

$(a2)$ *$E(A)$ is an $\sim$-equivalence class of $M_1(G)$ for every component A of G.*

Proof (uses Theorem 3.2.7 and 3.3.7 and Claim 4.3.3)

(**p1**) First we prove $(a1) \Rightarrow (a2)$. Let A be a component of G and $a, b \in E(G)$.

First suppose that a and b both belong to A. Since $G \in \mathcal{G}_{\bowtie}$, clearly also $A \in \mathcal{G}_{\bowtie}$. By Theorem 3.2.7, there exists a bicycle A_0 such that $A_0 \subseteq^s A$ and $a, b \in A_0$. Since A_0 is a bicycle, $A_0 \in \mathcal{C}_1(G)$, and therefore $a \sim b$.

Now suppose that a and b do not belong to the same component of G. Then no $C \in \mathcal{C}_1(G)$ contains both a and b because $G\langle C\rangle$ is a bicycle, which is a connected graph.

Therefore $E(A)$ is an $\sim$-equivalence class of $M_1(G)$.

(p2) Now we prove $(a2) \Rightarrow (a1)$. Let A be a component of G.

Since $E(A)$ is an $\sim$-equivalence class of $M_1(G)$, clearly $\mathcal{C}_1(G) \neq \emptyset$, and therefore matroid $M_k(G)$ is non-trivial and the core $[G]$ of G is defined.

Since $E(A)$ is an $\sim$-equivalence class of $M_1(G)$ for every component A of G, we also have: every component A has at least two elements that belong to a common circuit of $M_1(G)$. Therefore $L^*_k(G) = \emptyset$. Then by Claim 4.3.3, $E = E[G]$, and so $G = [G]$. Now by Theorem 3.3.7, $G \in \mathcal{G}_{\bowtie}$. $\square$

Corollary 4.4.7 *Let G be a graph. Suppose that $M_1(G)$ is a non-trivial matroid and $G \in \mathcal{G}_{\bowtie}$. Then N is a component of $M_1(G)$ if and only if $N = M_1(A)$ for some component A of G.*

Remark 4.4.8 Because of Corollary 4.4.7 we can reduce our problem of describing the classes of graphs having the same non-trivial bicircular matroid to the corresponding problem for the graphs in $\mathcal{CG}_{\bowtie}$, i.e. for connected cacti-graphs.

For this reasons from now on in the case of bicircular matroid we will assume that $G \in \mathcal{CG}_{\bowtie}$.

From Corollary 4.4.7 we have:

Theorem 4.4.9 GRAPH DESCRIPTION OF CONNECTED MATROID $M_1(G)$

The following are equivalent:

$(a1)$ *$M_1(G)$ is a non-trivial matroid and $G \in \mathcal{CG}_{\bowtie}$ and*

$(a2)$ *$M_1(G)$ is a connected matroid.*

From Theorems 4.2.1, 4.4.5, and 4.4.9 we have:

Claim 4.4.10 *Let G be a 2-connected graph and $k \geq 1$. If $\Delta G \geq k$, then $M_k(G)$ is a connected matroid.*

From Theorems 3.3.7, 4.2.1, 4.4.5, and 4.4.9 we also have:

Claim 4.4.11 *Let $k \geq 1$. If $M_k(G)$ is a connected matroid, then $G \in \mathcal{G}_{\bowtie}$ and $\Delta G \geq k$.*

Remark 4.4.12 The results in this section will allow us to reduce our problem $(WP)_k$ of describing the classes of graphs having the same k-circular matroid to the corresponding problem for cacti-graphs graphs, when $2 \leq k \leq \Delta G$, and for connected cacti-graphs, when $k = 1$. Equivalently, the theorems in this section allow us to reduce our problem $(WP)_k$ of finding all graph representations of a given k-circular matroid, $k \geq 1$, to the corresponding problem for the class of connected k-circular matroids.

4.5 Independent sets and bases of a k-circular matroid

Claim 4.5.1 *Let $G = (V, E, \phi)$ and $X \subseteq E$. If $\Delta G\langle X\rangle < k$ and $G\langle X\rangle$ has no tree component, then $X \in \mathcal{I}_k(G)$.*

Proof (uses Claim 3.1.5)

Let $X \subseteq E$ and $\Delta G\langle X\rangle < k$. If $G\langle X\rangle$ has no tree-component, then by Claim 3.1.5, $\Delta G\langle Z\rangle \leq \Delta G\langle X\rangle < k$ for every $Z \subseteq X$. Hence X contains no circuit of $M_k(G)$ and therefore X is an independent set of $M_k(G)$. ☐

Claim 4.5.2 *Let G be graph and $k \geq 1$. Suppose that $I \in \mathcal{I}_k(G)$, T is a tree component of $G\langle I\rangle$ or T is a vertex of G not in $G\langle I\rangle$, and e is an edge in $E(G) \setminus I$ incident to at least one vertex of T. Then $I \cup e \in \mathcal{I}_k(G)$.*

Proof (uses Claim 4.1.8)

Suppose, to the contrary, that $I \cup e \notin \mathcal{I}_k(G)$. Then there exists a unique $C \in \mathcal{C}_k(G)$ such that $e \in C \subseteq I \cup e$. If e has exactly one end-vertex in T, then the component of $G\langle C\rangle$ that contains e will contain a leaf, contradicting Claim 4.1.8. If both ends of e belong to T and T' is the graph obtained from T by adding edge e, then is connected and has at most one cycle. Since the component A of $G\langle C\rangle$ containing e is a subgraph of T', clearly A also has at most one cycle contradicting Claim 4.1.8 ($c2$). ☐

Theorem 4.5.3 GRAPH STRUCTURE OF A BASE OF $M_k(G)$ IN GRAPH G

Let G be a graph and $k \geq 1$. Suppose that $M_k(G)$ is a connected matroid. Then the following are equivalent:

($c1$) *$B \in \mathcal{B}_k(G)$ and*

($c2$) *$\Delta G\langle B\rangle = k - 1$, $V(G\langle B\rangle) = V(G)$ (i.e. B spans $V(G)$), and $\Delta A \geq 0$ for every component A of $G\langle B\rangle$ (i.e. $G\langle B\rangle$ has no tree component).*

Proof (uses Claims 3.1.7, 3.1.4, 3.1.5, 4.1.15, 4.3.2, 4.5.1, and 4.5.2)

Since $M_k(G)$ is a connected matroid, by Theorem 4.4.5 for $k \geq 2$ and Theorem 4.4.9 for $k = 1$, matroid $M_k(G)$ is non-trivial and also graph G is a cacti-graph. Therefore graph G has no tree component.

(**p1**) First we prove that ($c1$) $\Rightarrow$ ($c2$). Let $B \in \mathcal{B}_k(G)$.

(**p1.1**) Our first step is to show that $V(G) = V(G\langle B\rangle)$. Suppose, to the contrary, that there exists vertex v in $V(G) \setminus V(G\langle B\rangle)$. Since G has no isolated vertices, there exists $e \in E(G) \setminus B$ such that e is incident to v in G. Then by Lemma 4.5.2 (where T is vertex v), $B \cup e \in \mathcal{I}_k(G)$. This contradicts the maximality of B. Hence, if $B \in \mathcal{B}_k(G)$, then $V(G\langle B\rangle) = V(G)$.

(**p1.2**) Our next step is to show that $G\langle B\rangle$ has no tree component.

Suppose first that $G\langle B\rangle$ is a spanning tree. Since $M_k(G)$ is a non-trivial matroid, there exists $e \in E(G) \setminus B$. By maximality of B, we have: $B \cup e$ is a dependent set of $M_k(G)$.

Note that $\Delta_G(B \cup e) = 0$. Since $G\langle B \cup e\rangle$ is connected, by Claim 3.1.4. we have: $\Delta G\langle X\rangle \leq \Delta G\langle B \cup e\rangle = 0 < k$ for every $X \subseteq B \cup e$. Therefore $B \cup e$ contains no k-circuit, a contradiction.

Now suppose that $G\langle B\rangle$ has more than one component and one of them is a tree component, say T. Since G has no tree component, there exists $e \in E(G) \setminus B$ such that e is incident to at least one vertex of T. Then by Lemma 4.5.2, $B \cup e \in \mathcal{I}_k(G)$, a contradiction. Therefore $G\langle B\rangle$ has no tree component (i.e. $\Delta A \geq 0$ for every component A of $G\langle B\rangle$).

(**p1.3**) Now we show that $\Delta G\langle B\rangle = k-1$. Suppose, to the contrary, that $\Delta G\langle B\rangle \neq k-1$. By Claim 4.1.15, $\Delta G\langle B\rangle < k$. Therefore $\Delta G\langle B\rangle < k-1$. Since $M_k(G)$ is a non-trivial matroid, there exists edge e in $E(G) \setminus B$. By Claim 3.1.7, $\Delta G\langle B \cup e\rangle \leq k-1 < k$. Since $V(G\langle B\rangle) = V(G)$ and G has no tree component, clearly $G\langle B\rangle$ has no tree component as well. Now by Claim 4.5.1, $B \cup e \in \mathcal{I}_k(G)$. Therefore B is not a maximal independent set of $M_k(G)$, and so B is not a base of $M_k(G)$, a contradiction. Thus, $\Delta G\langle B\rangle \geq k-1$, that contradicts our assumption that $\Delta G\langle B\rangle < k-1$.

(**p2**) Finally, we prove $(c1) \Leftarrow (c2)$. Since no component of $G\langle B\rangle$ is a tree, by Claim 3.1.5, $\Delta G\langle X\rangle \leq \Delta G\langle B\rangle = k-1 < k$ for every $X \subseteq B$. Therefore B contains no circuit of $M_k(G)$, and so $B \in \mathcal{I}_k(G)$. Since $V(G\langle B\rangle = V(B)$, we have: $\Delta G\langle B \cup e\rangle > \Delta G\langle B\rangle = k-1$ for every $e \in E(G) \setminus B$. Then, by Claim 4.1.15, $B \cup e \in \mathcal{D}_k(G)$ for every $e \in E(G) \setminus B$. Therefore B is a maximal independent set of $M_k(G)$. $\square$

Let $\rho_k(G)$ and $\rho^*_k(G)$ denote the rank and the corank of matroid $M_k(G)$, respectively, and so $\rho_k(G) + \rho^*_k(G) = |E(G)|$. Obviously, if $\mathcal{C}_k(G) = \emptyset$, then $\rho_k(G) = |E(G)|$ and $\rho^*_k(G) = 0$. We also know that if G has no isolated vertices, then $\rho_0(G) = |V(G)| - cmp(G)$, where $cmp(G)$ is the number of components of G.

Corollary 4.5.4 *Let G be a graph and $k \geq 1$. Suppose that $M_k(G)$ is a connected matroid. Then $\rho_k(G) = |V(G)| - 1 + k$ and $\rho^*_k(G) = |E(G)| - |V(G)| + 1 - k$.*

Proof (uses Claim 4.5.3)
By Claim 4.5.3 $(c2)$, if B is a base of $M_k(G)$, then
$|B| - |V(G\langle B\rangle)| = \Delta G\langle B\rangle = k-1$ and $V(G\langle B\rangle) = V(G)$. Therefore
$\rho_k(G) = |B| = |V(G\langle B\rangle)| - 1 + k = |V(G)| - 1 + k$ and $\rho^*_k(G) = |E(G)| - |V(G)| + 1 - k$. $\square$

4.6 The core of a base of a k-circular matroid

In this Subsection we define the notion and describe some properties of the core of a base of $M_k(G)$ as a special subgraph of G. This notion will be essential for establishing some properties of the cocircuits of $M_k(G)$ in Subsection 4.7.

The definition below uses the notion of the core $[G]$ of a graph G (see Definition 3.3.4).

Definition 4.6.1 THE CORE OF A BASE OF M_k(G).
Suppose that $k \geq 2$, $M_k(G)$ is a connected matroid, and B is a base of $M_k(G)$.
Then the core $[G\langle B\rangle]$ of graph G is also called *the core of B in G*.

By Theorem 4.5.3, if $k \ge 2$ and B is a base of a connected matroid $M_k(G)$, then the core $[G\langle B\rangle]$ of graph $G\langle B\rangle$ is defined and if $k = 1$, then the core of B in G is not defined.

From Theorems 3.3.7 ($a3$) and 4.1.8 we have:

Claim 4.6.2 *Let $k \ge 2$ and $B \in \mathcal{B}_k(G)$. Suppose that $M_k(G)$ is a connected matroid. Then*

($c1$) *$[G\langle B\rangle]$ is the unique subgraph of $G\langle B\rangle$ belonging to $\mathcal{G}_{\bowtie}$ such that $\Delta[G\langle B\rangle] = \Delta G\langle B\rangle = k - 1$ and*

($c2$) *the edge set of $[G\langle B\rangle]$ is the unique subset of B that is a (k-1)-circuit of G.*

Claim 4.6.3 *Let $k \ge 2$ and $C \in \mathcal{C}_{k-1}(G)$. Suppose that $M_k(G)$ is a connected matroid. Then there exists $B \in \mathcal{B}_k(G)$ such that $G\langle C\rangle = [G\langle B\rangle]$.*

Proof (uses Theorem 4.5.3)

Let $Q \in Cmp(G)$. If $C \cap E(Q) \ne \emptyset$, then let Q' be a minimal connected spanning subgraph of Q containing C. If $C \cap E(Q) = \emptyset$, then let Q' be a minimal connected unicyclic subgraph of Q. Clearly, $H = \cup\{Q' : Q \in Cmp(G)\}$ satisfies condition ($c2$) of Theorem 4.5.3. Therefore $B = E(H) \in \mathcal{B}_k(G)$ and $G\langle C\rangle = [G\langle B\rangle]$. □

From Claims 4.6.2 and 4.6.3 we have:

Claim 4.6.4 *Let $k \ge 2$. Suppose that $M_k(G)$ is a connected matroid. Then $\mathcal{C}_{k-1}(G) = \{[G\langle B\rangle] : B \in \mathcal{B}_k(G)\}$.*

The claim below uses the notion of the kernel $\lfloor G\rfloor$ of a graph G (see Definition 3.3.3).

Claim 4.6.5 *Let $k = 1$. Suppose that $M_1(G)$ is a connected matroid. Then*

($c1$) *if $B \in \mathcal{B}_1(G)$, then $\lfloor A\rfloor$ is a cycle for every $A \in Cmp(G\langle B\rangle)$ and*

($c2$) *if C is the edge set of a cycle of G, then there exists $B \in \mathcal{B}_1(G)$ and $A \in Cmp(G\langle B\rangle)$ such that $G\langle C\rangle = \lfloor A\rfloor$.*

Proof (uses Theorem 4.5.3)

First we prove ($c1$).By Theorem 4.5.3, $\Delta G\langle B\rangle = 0$ and $\Delta A \ge 0$ for every $A \in Cmp(G\langle B\rangle)$. Now $G\langle B\rangle = \cup\{A : A \in Cmp(G\langle B\rangle)\}$, and so $\Delta G\langle B\rangle = \sum\{\Delta A : A \in Cmp(G\langle B\rangle)\}$. Therefore $\Delta A = 0$ for every $A \in Cmp(G\langle B\rangle)$. Then every component A of $G\langle B\rangle$. Thus $\lfloor A\rfloor$ is a cycle for every $A \in Cmp(G\langle B\rangle)$.

Now we prove ($c2$). Let C be a cycle of G. Let $A \in Cmp(G)$. If A contains C, then let A' be a minimal connected spanning subgraph of A containing C. If A does not contain C, then let A' be a minimal connected unicyclic spanning subgraph of A containing C. Clearly, $H = \cup\{A' : A \in Cmp(G)\}$ satisfies condition ($c2$) of Theorem 4.5.3. Therefore $B = E(H) \in \mathcal{B}_k(G)$ and $G\langle C\rangle = \lfloor A\rfloor$ for some $A \in Cmp(G)$. □

4.7 Cocircuits of a k-circular matroid

In this Section we intend to describe the cocircuits of the k-circular matroid M_k of a graph G as some special edge subsets of G. For example, we know that K is a cocircuit of $M_0(G)$ if and only if K is a minimal edge cut in G.

By Claim 2.2.4 ($c3$), for every cocircuit K of a matroid M there exists a base B of M and $e \in B$ such that K is the fundamental cocircuit of base B in M rooted at e (or the same, a (B, e)-cocircuit in M), i.e. $K = C^*(e, B)$ for some $B \in \mathcal{B}(M)$ and $e \in B$. Put $K(e, B) = C^*(e, B)$.

Thus, we can (and will) describe all cocircuits of matroid $M_k(G)$ by describing the (B, e)-cocircuits $K(e, B)$ of $M_k(G)$ for all pairs (B, e), where B is a base of $M_k(G)$ and $e \in B$.

We will distinguish between three possible types of (B, e)-cocircuits $K(e, B)$ depending on the structure of component A of $G\langle B\rangle$ containing edge e and on the position of edge e in A.

Definition 4.7.1 *Let $B \in \mathcal{B}_k(G)$, $e \in B$, and $k \geq 1$. Then*

($t1$) *$K(e, B)$ is a (B, e)-*cocircuit in $M_k(G)$ of type 1*, if $e \notin E\lfloor A\rfloor$, where A is a component of $G\langle B\rangle$ containing edge e,*

($t2$) *$K(e, B)$ is a (B, e)-*cocircuit in $M_k(G)$ of type 2*, if $e \in E\lfloor A\rfloor$, where A is a unicyclic component of $G\langle B\rangle$ containing edge e, and*

($t3$) *$K(e, B)$ is a (B, e)-*cocircuit in $M_k(G)$ of type 3*, if $e \in E[A]$, where A is a component of $G\langle B\rangle$ that has at least two cycles and contains edge e.*

Theorem 4.7.2 GRAPH DESCRIPTION OF ROOTED COCIRCUITS OF TYPE 1

Let G be a graph and $k \geq 1$. Suppose that

($a1$) *$M_k(G)$ is a connected matroid, $B \in \mathcal{B}_k(G)$, and $e \in B$ and*

($a2$) *$K(e, B)$ is a (B, e)-cocircuit in $M_k(G)$ of type 1 (i.e. edge $e \notin E\lfloor A\rfloor$, where A is a component of $G\langle B\rangle$ containing edge e).*

Then exactly one of the two components of $A \setminus e$ is a tree T and $K(e, B) = K'(e, B) \cup e$, where $K'(e, B)$ is the set of edges in $E \setminus B$ having at least one end-vertex in $V(T)$.

Proof (uses Theorem 4.5.3 and Claims 2.2.4 and 3.3.9)

(**p1**) Since $e \notin E\lfloor A\rfloor$, by Claim 3.3.9 ($c1$), exactly one of the two components of $A \setminus e$ is a tree. Let T be the tree component and D the non-tree component of $A \setminus e$.

(**p2**) We prove that $K(e, B) = K'(e, B) \cup e$. By Claim 2.2.4 ($c2$), it is sufficient to show that $u \in K'(e, B)$ if and only if $u \in E \setminus B$ and $B_u = (B \setminus e) \cup u$ satisfies condition ($c2$) of Theorem 4.5.3.

(**p2.1**) First, suppose that $u \in K'(e, B)$.

If u has both ends in $V(T)$, then both D and $T \cup u$ have cycles, and therefore B_u satisfies condition ($c2$) of Theorem 4.5.3.

If u has one end in T and the other end in D, then $G\langle(E(A) \setminus e) \cup u\rangle$ is not a tree and B_u satisfies condition ($c2$) of Theorem 4.5.3.

If u has exactly one end in T and the other end in a component F of $G\langle B\rangle$ distinct from A, then $G\langle E(F) \cup E(T) \cup u\rangle$ is not a tree, and therefore B_u satisfies condition $(c2)$ of Theorem 4.5.3.

(**p2.2**) Now suppose that $u \in E \setminus K(e, B)$. Then T is a tree-component of $G\langle B \setminus e \cup u\rangle$, and so B_u does not satisfies condition $(c2)$ of Claim 4.5.3. $\square$

Theorem 4.7.3 GRAPH DESCRIPTION OF ROOTED COCIRCUITS OF TYPE 2

Let G be a graph and $k \geq 1$. Suppose that

$(a1)$ *$M_k(G)$ is a connected matroid, $B \in \mathcal{B}_k(G)$, and $e \in B$ and*

$(a2)$ *$K(e, B)$ is a (B, e)-cocircuit in $M_k(G)$ of type 2, (i.e. $e \in E\lfloor A\rfloor$, where A is a unicyclic component of $G\langle B\rangle$ containing edge e).*

Then $A \setminus e$ is a tree and $K(e, B) = K'(e, B) \cup e$, where $K'(e, B)$ is the set of edges in $E \setminus B$ having at least one end-vertex in $V(A \setminus e)$.

Proof (uses Theorem 4.5.3 and Claim 2.2.4)

By assumption $(a2)$ of our theorem, e is an edge of a unique cycle in A. If u has both ends in $V(A)$, then $\langle (E(A) \setminus e) \cup u\rangle$ is not a tree and therefore $(B \setminus e) \cup u$ satisfies condition $(c2)$ of Theorem 4.5.3.

If u has exactly one end in $V(A)$ and the other end in a component D of $G\langle B \setminus e\rangle$, then $G\langle E(D) \cup E(A \setminus e) \cup u\rangle$ is not a tree and $(B \setminus e) \cup u$ satisfies condition $(c2)$ of Theorem 4.5.3.

Now, if u has no ends in $E(A)$ or, equivalently, $u \notin K'(e, B)$, then $A \setminus e$ is a tree component of $G\langle (B \setminus e) \cup u\rangle$, and so $(B \setminus e) \cup u$ does not satisfies condition $(c2)$ of Theorem 4.5.3. Therefore $(B \setminus e) \cup u \notin \mathcal{B}_k(G)$.

Hence by Claim 2.2.4 $(c2)$,
$u \in K'(e, B) \cup e = K(e, B) \Leftrightarrow (B \setminus e) \cup u \in \mathcal{B}_k(G) \Leftrightarrow u \in K(e, B)$. $\square$

Theorem 4.7.4 GRAPH DESCRIPTION OF ROOTED COCIRCUITS OF TYPE 3

Let G be a graph and $k \geq 1$. Suppose that

$(a1)$ *$M_k(G)$ is a connected matroid, $B \in \mathcal{B}_k(G)$, and $e \in B$ and*

$(a2)$ *$K(e, B)$ is a (B, e)-cocircuit in $M_k(G)$ of type 3, (i.e. $e \in E[A]$, where A is a component of $G\langle B\rangle$ that has at least two cycles and contains edge e).*

Then $K(e, B) = (E \setminus B) \cup e$.

Proof (uses Theorem 4.5.3 and Claim 3.3.9)

Since $e \in E[A]$, by Claim 3.3.9 $(c3)$, every component of $A \setminus e$ contains a cycle. Therefore for every $u \in E \setminus B$ we have: $(B \setminus e) \cup u$ satisfies condition $(c2)$ of Theorem 4.5.3. Hence $u \in K(e, B)$ if and only if $u \in (E \setminus B) \cup e$. $\square$

References

[1] J. A. Bondy and U. S. R. Murty, *Graph Theory*, 3rd Corrected Printing, GTM 244, Springer-Verlag, New York, 2008.

[2] C. R. Coulard, J. G. del Greco, D. K. Wagner, Representations of bicircular matroids, *Discrete Applied Mathematics* **32** (1991) 223-240.

[3] R. Diestel, *Graph Theory*, Springer-Verlag, New York, 2000.

[4] J. Edmonds and G.-C. Rota, Submodular set functions (Abstract), *Waterloo combinatorics conference*, (1966).

[5] R. Halin and H. A. Jung, Note on isomorphisms of graphs, *J. London Math.* Sec. **42** (1967) 254-256.

[6] R. L. Hemminger, H. A. Jung, A. K. Kelmans, On 3–skein isomorphisms of a graph, *Combinatorica* **2** (4), (1982) 373-376.

[7] A. K. Kelmans, On semi-isomorphisms and semi-dualities of graphs, *Graphs and Combinatorics* **10** (1994) 337–352.

[8] A. K. Kelmans, Matroids and combinatorial optimization problems. Institute of Control Sciences, Moscow (1989).

[9] A. K. Kelmans, On 3-skeins in a 3-connected graph. *Studia Scientarum Mathematicarum Hungarica* **22** (1987) 265-273.

[10] J. G. Oxley, *Matroid Theory*, Oxford University Press, 2006.

[11] D. K. Wagner, Connectivity in Bicircular Matroids, *J. Combinatorial Theory* **B 39** (1985) 308-324.

[12] D. J. A. Welsh, *Matroid Theory*, Academic Press, London, 1976.

[13] D. West, *Introduction to Graph Theory*, Prentice Hall, 2001.

[14] H. Whitney, On the abstract properties of linear dependence, *Amer. J. Math.* **57** (1935) 509-533.

[15] H. Whitney, 2-isomorphic graphs, *Amer. J. Math.* **55** (1933) 245-254.

[16] H. Whitney, Congruent graphs and the connectivity of graphs, *Amer. J. Math.* **54** (1932) 159-168.

[17] H. Whitney, Non-separable and planar graphs, *Amer. J. Math.* **34** (1932) 339-362.